

# 1 Scattering by nonspherical particles (lecture 14)

Perfectly spherical particles constitute, practically, an exception in nature and even in industrial applications. In the recent past, numerical methods have been actively developed for light scattering by nonspherical particles. In practice, the methods require extensive computational capacity including supercomputers.

In what follows, one possible modeling of a nonspherical particle geometry is presented: the Gaussian random sphere. Thereafter, computation of scattering by Gaussian particles is discussed in various approximations, whereafter a summary is given on essentially exact numerical methods and possibilities to apply these methods to scattering by Gaussian particles.

## 2 Gaussian random particle

Statistical modeling of nonspherical particle shapes seems reasonable, since nonspherical shapes usually show up as a wide spectrum of different-looking shapes. In the Gaussian-random-sphere model, the particle is assumed to be mathematically star-like so that there is an origin with respect to which the shape can be expressed as a function of the spherical coordinates. In the spherical geometry, the so-called lognormal statistics are being used so that the radial distance of the particle varies within  $]0, \infty[$ . The shape is unambiguously defined by the mean of the radial distance  $a$  and the covariance function of the logarithm of the radial distance  $\Sigma_s$ . Explicitly,

$$r(\theta, \varphi) = ae^{s(\theta, \varphi) - \frac{1}{2}\beta^2},$$

where  $s$  is the logarithmic radial distance and  $\beta^2 = \Sigma_s(0)$  is the variance of  $s$ . Now

$$s(\theta, \varphi) = \sum_{lm} s_{lm} Y_{lm}(\theta, \varphi)$$

and, due to  $s$  being real-valued,

$$s_{l,-m} = (-1)^m s_{lm}^* \begin{cases} l = 0, 1, 2, \dots, & \\ m = -l, \dots, -1, 0, 1, \dots, l, & \end{cases} ;$$

$$Im(s_{l0}) = 0.$$

The spherical harmonics coefficients of the logarithmic radial distance  $s_{lm}, m \geq 0$  are independent Gaussian random variables with zero means and with variances ( $l$  and  $m$  as above)

$$\begin{aligned} \text{Var}[\Re(s_{lm})] &= (1 + \delta_{m0}) \frac{2\pi}{2l+1} c_l \\ \text{Var}[\Im(s_{lm})] &= (1 - \delta_{m0}) \frac{2\pi}{2l+1} c_l \end{aligned}$$

The coefficients  $c_l \geq 0, l = 0, \dots, \infty$  are the coefficients of the Legendre expansion for the covariance function  $\Sigma_s$ :

$$\Sigma_s(\gamma) = \beta^2 C_s(\gamma) = \sum_{l=0}^{\infty} c_l P_l(\cos \gamma), \quad \sum_{l=0}^{\infty} c_l = \beta^2,$$

where  $\gamma$  is the angular distance between two directions  $(\theta_1, \varphi_1)$  and  $(\theta_2, \varphi_2)$ .

The two slopes on the Gaussian random particle (subscripts referring to partial derivatives)

$$s_\theta = \frac{r_\theta}{r}, \quad \frac{1}{\sin \theta} s_\varphi = \frac{r_\varphi}{r \sin \theta}$$

are, again, independent Gaussian random variables with zero means and with standard deviations

$$\rho = \sqrt{-\Sigma_s^{(2)}(0)},$$

where  $\Sigma_s^{(2)}$  is the second derivative of the covariance function with respect to  $\gamma$ . The correlation length  $l_c$  and correlation angle  $\Gamma_c$  are

$$l_c = 2 \sin \frac{1}{2} \Gamma_c = \frac{1}{\sqrt{-c_s^{(2)}(0)}}.$$

Natural random shapes often exhibit covariance functions, for which the coefficients  $c_l$  follow the exponent form  $c_l \propto l^{-\nu}$ ,  $l \geq 2$ . For  $\nu = 4$ , one obtains random shapes applicable, in the first place, to modeling Saharan sand particles, asteroids, as well as the shapes of terrestrial planets.

In the limiting case, the Gaussian random shape thus depends on a single free parameter insofar as the shape is concerned: the variance  $\beta^2$  of the logarithmic radial distance.  $\beta^2$  relates to the relative variance of the radius  $\sigma^2$  via the simple relation

$$\sigma^2 = e^{\beta^2} - 1.$$

Increasing  $\sigma$  results in shapes, where the radial fluctuations are enhanced.

If, additionally,  $\nu$  is treated as a free parameter, one obtains shorter correlation lengths with smaller values of  $\nu$  (when the expansions are always truncated at a certain degree  $l_{max}$ ) and thereby larger numbers of hills and valleys as per unit solid angle.

For  $\nu \geq 4$ , non-fractal smooth shapes are obtained whereas, for  $\nu < 4$ , fractal shapes follow, in which case infinite expansions would yield non-differentiable surfaces rendering the discussion of slopes meaningless.

### 3 Scattering by Gaussian particles in different approximations

Light scattering by Gaussian random particles has been studied in the ray-optics, Rayleigh-volume, Rayleigh-Gans, anomalous-diffraction and perturbation-series approximations, as well as in the Rayleigh-ellipsoid approximation.

In the Rayleigh-volume approximation, the scattering by a small particle follows from its volume. In the case of the Gaussian particle, the (ensemble-averaged) absorption cross section

is proportional to the mean of the volume, whereas the scattering cross section is proportional to the mean of the squared volume. The angular characteristics of the scattering matrix are the same as in the Rayleigh approximation for spherical particles. The results are largely analytical.

In the Rayleigh-ellipsoid approximation, an ellipsoid is fitted to each realization of the Gaussian particle, the ellipsoid volume being equal to the volume of the realization. Scattering is then approximated with the existing electrostatics approximation for ellipsoidal scatterers. The most significant challenge in the Rayleigh-ellipsoid approximation is the numerical computation of the best-fit ellipsoid, whereafter the results follow in a straightforward way.

In the Rayleigh-Gans approximation (or the first Born approximation), the numerical computation of the form factor can be aided by analytical intermediate results. In practice, some numerical integration remains, preventing the treatment of arbitrarily large particles.

In anomalous diffraction, path lengths of rays inside the Gaussian sample particles are numerically computed in cases where the refractive index is close to unity. The absorption follows directly from the exponential attenuation and extinction is computed from the optical theorem. The angular dependence of scattering is obtained by averaging the square of the scattering amplitude. The most demanding task is the computation of the path lengths inside the particle, which is difficult for extremely nonspherical shapes.

In the second-order perturbation-series approach for the boundary conditions, analytical results follow for the cross sections and scattering matrices and the most challenging numerical part is the computation of the so-called  $3j$ -symbols. The unknown accuracy of the results is a problem. In practice, the perturbation-series method is applicable to wavelength-scale scatterers only, if the deviations from the spherical shape are small compared to the wavelength.

Approximations can be taken to be "the spice" that makes the scattering research "delicious", since, in practice, all so-called exact methods are based on approximation in some part. One can make the provocative statement that only approximations allow the computation of light scattering by realistic small particles. The applicability of the exact methods is usually limited to a narrow range of simple shapes. By the rapid development of computers and by the development of new analytical methods, the applicability of certain exact methods grows slowly but steadily.

## 4 Exact methods and their applicability to Gaussian particles

The numerical methods in light scattering can be divided into differential-equation and integral-equation methods. The traditional computational method is the separation-of-variables method that has been successful in the solution of the following scattering problems:

1. isotropic, homogeneous sphere

2. coated sphere consisting of the interior and coating (with common origin)
3. layered sphere that consists of several layers defined by concentric spherical cells
4. radially inhomogeneous sphere
5. optically active (chiral) sphere
6. homogeneous, isotropic infinite circular cylinder
7. optically active infinite circular cylinder
8. isotropic infinite elliptic cylinder
9. isotropic, homogeneous spheroid
10. coated spheroid that consists of the interior and coating (with common origin)
11. optically active spheroid

The separation-of-variables method is not applicable to scattering by Gaussian particles.

The FEM-method (finite-element method) is a differential-equation method, where the scatterer is placed in a finite computational volume that is discretized into numerous small computational cells. Typically, there are 10-20 cells per wavelength and the electromagnetic field is solved for in the nodal points of the cells. The resulting linear group of equations consists of a sparse matrix. In the boundaries of the computational volume, an artificial absorbing boundary condition is invoked. Although FEM allows for the computations for arbitrary, even inhomogeneous particles, it has not yet been applied to Gaussian particles.

The FDTD-method (finite-difference time-domain method) is a differential-equation method that solves for the time dependence of the electromagnetic fields based on Maxwell's curl equations. Both time and spatial derivatives are expressed with finite differences and time elapses in finite steps. The scattering particle is again placed in a finite computational volume and an absorbing boundary condition is required in the boundary of the computational volume. The density of the discretization is as in the FEM-method. In FDTD, there is no need to solve a large group of equations. Recently, the method has yielded promising results in light scattering by Gaussian particles.

In the PM-method (point matching), the boundary conditions of the electromagnetic fields are required in a finite number of points on the surface of the particle. In the original method, there were as many points as unknown coefficients in the vector spherical harmonics expansion. It was concluded that the method was numerically unstable. There is, however, nothing that prevents us from expanding the number of points and computing the coefficients using the least-squares method. This version of the method has been noticed to be stable and is one of the most popular numerical methods. The regime of application can be improved by expanding the fields with a number of suitably chosen origins within the particle. PM is promising also

for scattering by Gaussian particles. It is intriguing to ponder whether “an educated guess” can help speed up the solution of the coefficients.

The integral-equation methods are divided into a wide spectrum of different methods. In the VIEM method (volume-integral-equation), one considers the integral equation

$$\mathbf{E}(\mathbf{r}) = \mathbf{E}_i(\mathbf{r}) + k^2 \int_V d^3\mathbf{r}' \left[ \mathbf{1} + \frac{1}{k^2} \nabla \nabla \right] \frac{e^{ik|\mathbf{r}-\mathbf{r}'|}}{4\pi|\mathbf{r}-\mathbf{r}'|} \cdot [m^2(\mathbf{r}') - 1] \mathbf{E}(\mathbf{r}').$$

By discretizing the integral on the right-hand side, one obtains a group of linear equations for the field values at the discretization points within the volume of the particle. Solving the equations results in the field inside the particle. Typically, again, 10-20 discretization points are required as per wavelength so that, after a straightforward calculation, it is clear that a group of equations with thousands of unknowns easily follows. In practice with current computers, up to 200 million unknowns can be treated (as of December 12, 2008). Various versions of the VIEM method have been successfully applied to Gaussian-particle scattering (foremost DDA, discrete-dipole approximation).

In the case of VIEM, the matrix of the group of linear equations is full, which makes the solution more difficult. When the internal field has been solved for, the same integral relation gives the scattered field outside the particle via straightforward integration (subtracting the original field).

DDA (discrete-dipole approximation) is a certain version of solution methods for the integral equation. DDA can be visualized in the following: the particle can be thought to be composed of dipole scatterers interacting with each other. In practice, the VIEM methods differ from one another in how they treat the singular self-term inside the integral, which is essential for the accuracy of the method.

The surface-integral-equation methods (SIEM) make use of two-dimensional integral equations that seem like a reasonable starting point, in particular, for homogeneous particles. However, the SIEM-methods are less stable than the VIEM-methods and usually require additional regularization.

The integral equation shown above in connection to the VIEM-method is Fredholm-type and the kernel has a singularity at  $\mathbf{r} = \mathbf{r}'$ . Via Fourier-transformation, handling of the singularity can be improved and the integral equation can be solved numerically in the wavenumber (or frequency) space. Surprisingly, the disadvantage of the method is the considerable analytical work needed for each different particle. These so-called FIEM-methods have not been very popular.

In the TMM method (transition matrix method), the analysis proceeds with the help of vector spherical harmonics functions and the word “transition” refers to the linear matrix relation between the original field and the scattered field. Compared to the direct vector spherical harmonics treatment of the boundary conditions, TMM has the advantage that a linear relation is obtained purely between the internal and original fields, reducing the number of unknowns in the group of linear equations. After solving the group of equations,

the scattered is computed from the vector Kirchhoff integral relation. The TMM method is an efficient method, in particular, for axially symmetric particles and useful results have been obtained, e.g., for spheroids to compare with the implications of the SVM method. However, TMM suffers from unpredictable convergence and instability problems and have not yet been extensively applied to scattering by Gaussian particles. As a tool the actual  $T$ -matrix is quite useful and, for a single particle, needs to be computed only once (independently of the orientation). Recently, an analytical version of the  $T$ -matrix method has been developed—this version is highly promising for studying scattering by Gaussian random particles.

In the superposition method for spheres and spheroids, scattering by particle clusters is computed using the translation and addition rules of vector spherical harmonics functions. The field scattered by the cluster is expressed as a superposition of the fields scattered by each constituent particle. The partial fields depend on each other due to the mutual electromagnetic interactions of the constituent particles. The scattering problem again manifests itself in a solution of a group of linear equations. Currently, precise solutions can be computed for clusters with several dozens of constituent particles, when constituent-particle size approaches the wavelength.

## 5 Applications of electromagnetic scattering

In his book, van de Hulst has presented an excellent review of the applications of light scattering in various fields of science. This is recommended reading bearing in mind, in particular, modern computational methods for nonspherical particles. Bohren and Huffman offer additional material on the applications, as well as Mishchenko et al. Finally, the publications from the meeting series entitled *Electromagnetic and Light Scattering by Nonspherical Particles: Theory, Measurements, and Applications* offer up-to-date information about the advances in light scattering by small particles.