

Introduction to Conformal Geometry and Quasiconformal Maps
Department of Mathematics and Statistics
University of Helsinki
Winter 2011 / Vuorinen

Exercise 5, 2011-02-21 File: icg1105.tex, 2011-2-7,9.32

1. Let $G = \mathbb{R}^n \setminus \{0\}$, $x, y \in G$, and let $\varphi \in [0, \pi]$ be the angle between the segments $[0, x]$ and $[0, y]$.

(a) Show that $\sin \frac{1}{2}\varphi \leq \frac{|x-y|}{|x|+|y|}$.

(b) Show that $|x - y| \leq ||x| - |y|| + 2 \min\{|x|, |y|\} \sin(\varphi/2)$.

(c) It is known (by [MOS]) that $k_G(x, y) = \sqrt{\varphi^2 + \log^2 \frac{|x|}{|y|}}$. Using this result show that there is constant A such that that $k_G(x, y) \leq A j_G(x, y)$ for all $x, y \in G$ i.e. that G is a uniform domain.

2. Let $x, y \in \mathbb{R}^n \setminus \{0\}$ and $|y| \geq |x|$. Show that $d(y, [0, x]) \geq \frac{|x-y|}{2}$.

3. Let $f \in \mathcal{GM}(\mathbb{B}^n)$ and $r \in (0, 1)$. Show that

$$|f(x) - f(y)| \leq \frac{1}{1-r^2}|x - y|,$$

for $|x|, |y| \leq r$. [Hint: $\text{sh}^2 \frac{\rho(x,y)}{2} = \dots$]

4. For an open set D in \mathbb{R}^n , $D \neq \mathbb{R}^n$, let

$$\varphi_D(x, y) = \log \left(1 + \max \left\{ \frac{|x-y|}{\sqrt{d(x)d(y)}}, \frac{|x-y|^2}{d(x)d(y)} \right\} \right); \quad x, y \in D.$$

Show that $j_D(x, y)/2 \leq \varphi_D(x, y) \leq 2j_D(x, y)$.

5. Let $G = \mathbb{R}^n \setminus \{0\}$ and $f(x) = a^2x/|x|^2$ for $x \in G$, where $a > 0$. Show that $k_G(f(x), f(y)) = k_G(x, y)$ and $j_G(f(x), f(y)) = j_G(x, y)$ for $x, y \in G$.

6. Let $f : G \rightarrow G' = f(G)$, $G, G' \subset \mathbb{R}^n$, be a homeomorphism such that for some $C > 0$ and all $x, y \in G$, $k_{G'}(f(x), f(y)) \leq Ck_G(x, y)$. Suppose that $b \in \partial G$ and that $b_k \in G$ with $b_k \rightarrow b$, $f(b_k) \rightarrow \beta$, $k \rightarrow \infty$, and let $E = \cup D(b_k, 1)$. Here $D(x, M)$ stands for the quasihyperbolic ball. Prove that $f(x) \rightarrow \beta$ when $x \rightarrow b$, $x \in E$. Note: By topology, for each sequence (b_k) tending to a boundary point b of G such that the image sequence also has a limit γ , it follows that $\gamma \in \partial G'$.

File: icg1105.tex, 2011-2-7,9.32