

5th exercises for SIM'2020

Ex. 1

Load observations from `linear-model-data-3.dat`. First two columns are explanatory variables x_1, x_2 , third is dependent variable y . Find suitable model between y and x_i . You can test functions of x_i 's and interaction. Test at least two different models and choose best according to model selection criterion of your choice. Check if parameters in the model are significant and that residuals against predicted \hat{y} seem unbiased and homoscedastic. You can use ready-made regression package or write your own procedures.

Ex. 2

Fit nonlinear model to observed degree of linear polarization of comet Hale-Bopp in red wavelength filter. Download data from course webpage (`dataRed.dat`). Use trigonometric model

$$y_i = f_i(\boldsymbol{\beta}) = \beta_1 \sin(x_i)^{\beta_2} \cos(x_i/2)^{\beta_3} \sin(x_i - \beta_4). \quad (1)$$

- Program function $S(b_1, b_2, b_3, b_4) = \sum^n (y_i - f_i(b_1, b_2, b_3, b_4))^2$ in your computing environment. Use minimization procedure to find best estimates b_1, b_2, b_3, b_4 . (If you cannot use minimization, test few choices for parameters yourself and choose the best ones).
- Plot data together with the best fit function.
- Compute sum of squared residuals, SSE , and residual variance s^2 .
- Compute test statistics $t_i = \frac{b_i}{s\sqrt{m^{ii}}}$ for the parameters. For $m^{ii} = [M^{-1}]_{ii}$ you need the matrix $\mathbf{F}(\mathbf{b})$ as in Eq. (4.15). Partial derivatives of f are given in the end of this page.
- Which parameter is the most uncertain, i.e. has smallest value of test statistics? Test it's p -value for the hypothesis H_0 that it could be removed from the model.

$$\frac{\partial f}{\partial b_1} = -\sin(x)^{b_2} \cos\left(\frac{x}{2}\right)^{b_3} \sin(b_4 - x) \quad (2)$$

$$\frac{\partial f}{\partial b_2} = -b_1 \sin(x)^{b_2} \cos\left(\frac{x}{2}\right)^{b_3} \sin(b_4 - x) \log(\sin(x)) \quad (3)$$

$$\frac{\partial f}{\partial b_3} = -b_1 \sin(x)^{b_2} \cos\left(\frac{x}{2}\right)^{b_3} \sin(b_4 - x) \log\left(\cos\left(\frac{x}{2}\right)\right) \quad (4)$$

$$\frac{\partial f}{\partial b_4} = -b_1 \sin(x)^{b_2} \cos\left(\frac{x}{2}\right)^{b_3} \cos(b_4 - x) \quad (5)$$