

Computational light scattering, fall 2022 (PAP315, 5 cr), Exercise 2

The answers are due on **September 21, 2022**. Please return them to Anne Virkki via e-mail (anne.virkki@helsinki.fi).

1. Consider two small non-interacting spherical particles of radius $a \ll \lambda$, where λ is the wavelength of the incident plane wave (size parameter $x = 2\pi a/\lambda$). One particle is set to the origin and the location of the other particle is denoted by a vector \mathbf{d} . Study the interference in first-order scattering when the internal fields are assumed to coincide with the incident field at the particle locations.

(6 points)

2-3. A plane wave is scattered by two small interacting spherical particles of radius $a \ll \lambda$, where λ is the wavelength of the incident plane wave (size parameter $x = 2\pi a/\lambda$). One particle is set to the origin and the location of the other particle is denoted by a vector \mathbf{d} . In the dipole approximation, the internal fields of the particles \mathbf{E}_1 and \mathbf{E}_2 are related through

$$\begin{aligned}\mathbf{E}_1 &= \mathbf{E}_{i1} + \beta \bar{\mathbf{T}}(u, v) \cdot \mathbf{E}_2 \\ \mathbf{E}_2 &= \mathbf{E}_{i2} + \beta \bar{\mathbf{T}}(u, v) \cdot \mathbf{E}_1,\end{aligned}$$

where \mathbf{E}_{i1} and \mathbf{E}_{i2} are the incident fields at the locations of the particles, and the polarizability (m is the refractive index)

$$\beta = x^3 \frac{m^2 - 1}{m^2 + 2}.$$

The transformation $\bar{\mathbf{T}}$ denotes the interaction between the particles:

$$\begin{aligned}\bar{\mathbf{T}}(u, v) &= u\bar{\mathbf{I}} + v\mathbf{d}\mathbf{d}/d^2 \\ u &= e^{i\rho}(\rho^2 + i\rho - 1)/\rho^3 \\ v &= e^{i\rho}(-\rho^2 - i3\rho + 3)/\rho^3, \quad \rho = kd.\end{aligned}$$

Solve the electric fields \mathbf{E}_1 and \mathbf{E}_2 . (See Muinonen 1990, PhD thesis.)

Note the following rules for an operator \mathbf{ab} :

$$(\mathbf{ab}) \cdot \mathbf{c} = \mathbf{a}(\mathbf{b} \cdot \mathbf{c})$$

$$\mathbf{c} \cdot (\mathbf{ab}) = (\mathbf{c} \cdot \mathbf{a})\mathbf{b}$$

The unit operator $\bar{\mathbf{I}}$ has no effect on the operator \mathbf{ab} or the vector \mathbf{c} .

(12 points)

4. Calculate the form factor of the Rayleigh-Gans approximation for a cubic particle.

(6 points)