

Dependence logic  
 Problems 3  
 Tuesday 12.4.2011

1. Let  $\phi(x_1, \dots, x_n)$  and  $\psi(x_1, \dots, x_n)$  be formulas of dependence logic, with  $x_1, \dots, x_n$  appearing free, such that  $\phi(x_1, \dots, x_n) \equiv^* \psi(x_1, \dots, x_n)$ . For terms  $t_1, \dots, t_n$  let  $\phi(t_1, \dots, t_n)$  be the formula in which  $x_i$  is substituted by  $t_i$  for  $1 \leq i \leq n$ . The substitution is allowed only if no variable of  $t_i$  becomes bound in  $\phi(t_1, \dots, t_n)$ . Show that for any terms  $t_1, \dots, t_n$  satisfying this condition for both  $\phi$  and  $\psi$ ,

$$\phi(t_1, \dots, t_n) \equiv^* \psi(t_1, \dots, t_n).$$

2. Show the following implication for all sentences  $\phi$ , and models  $\mathcal{M}$  and  $\mathcal{M}'$ : if  $\mathcal{M} \cong \mathcal{M}'$ , then  $\mathcal{M} \models \phi \Leftrightarrow \mathcal{M}' \models \phi$ . Prove the claim using induction on  $\phi$  for all formulas. Consider the cases where  $\phi$  is an atomic formula or  $\phi = \exists x_n \psi$ .

3. Find a logically equivalent first-order formula in each case (below  $f$  and  $P$  are symbols of the vocabulary):

1.  $\exists x_0 (= (x_1, x_0) \wedge (f(x_1) = x_1))$
2.  $\exists x_0 (= (x_2, x_0) \wedge (P(f(x_0)) \wedge \neg P(x_1)))$

4. Which of the following formulas are logically equivalent to a first-order formula:

1.  $= (x_0, x_1, x_2) \wedge x_0 = x_1$
2.  $(= (x_0, x_2) \wedge x_0 = x_1) \rightarrow = (x_1, x_2)$
3.  $\forall x_0 \exists x_2 (= (x_0, x_2) \wedge x_2 = x_1)$

5. Let  $L = \emptyset$  and  $M = \{0, 1\}$ . Show that the following types of a team  $X$  of  $M$  with domain  $\{x_0, x_1, x_2\}$  are not first-order:

1.  $\exists x_0 (= (x_2, x_0) \wedge \neg (x_0 = x_1))$
2.  $\exists x_0 (= (x_2, x_0) \wedge (x_0 = x_1 \vee x_0 = x_2))$

6. Let  $\phi$  be the formula  $\exists x_0 \forall x_1 \neg (= (x_2, x_1) \wedge (x_0 = x_1))$ . Show that the flattening of  $\phi$  is not a strong logical consequence of  $\phi$ .