

Introduction to Conformal Geometry and Quasiconformal Maps
Department of Mathematics and Statistics
Winter 2013 / Vuorinen

Exercise 11, 2013 File: icg1311.tex, 2013-3-3,18.22

1. Let $c \geq 1$, and let $G \subset \mathbf{R}^n$ be an open set. A positive, continuous function $u : G \rightarrow R_+ \setminus \{0\}$ is called a c -Harnack function if the inequality

$$\sup_{\mathbf{B}(x,r)} u(z) \leq c \inf_{\mathbf{B}(x,r)} u(z)$$

holds whenever $\mathbf{B}(x, 2r) \subset G$. Well known examples of functions satisfying Harnack's inequality are positive harmonic functions in the plane.

(a) Let $u(z) = \arg z$ and $G = \mathbf{C} \setminus \{x \in \mathbf{R} : x \geq 0\}$. Find a constant $c \geq 1$ such that $u(z)$ is c -Harnack in G .

(b) Let $K \subset G$ be compact and $u(z)$ as in (a). Does there exist a constant D depending on $d(K)/d(K, \partial G)$ such that

$$u(z_1) \leq D u(z_2)$$

for all $z_1, z_2 \in K$?

(c) Let $K \subset G$ be compact. Show that $u(x) = \exp(-k_G(x, K))$ satisfies the Harnack inequality.

2. Let $G, G' \subset \mathbf{R}^n$ and $f : (G, k_G) \rightarrow (G', k'_G)$ be uniformly continuous, and let $b' \in \partial G'$. Show that $u : G \rightarrow R_+, u(x) = |f(x) - b'|$ satisfies Harnack's inequality.

3. Let $E \subset \mathbf{B}$ be compact. Suppose that

$$m_n(E_k) = a_k, E_k = \{x \in \mathbf{R}^n \setminus E : 2^{-k-1} < d(x, E) < 2^{-k}\}, k = 1, 2, \dots$$

Use Lemma 5.24[CGQM] to find an upper bound for $M(\Delta(E, S^{n-1}(2)))$. Apply your bound to give a sufficient condition for $\text{cap } E = 0$ in terms of the numbers (a_k) .

4. Let $G = \mathbf{B} \setminus \{0\}$, $f : G \rightarrow G' = f(G)$, be a homeomorphism with the property that there exist curves $\alpha_j : [0, 1) \rightarrow G, j = 1, 2$, such that $\alpha_j(t) \rightarrow 0, f(\alpha_j(t)) \rightarrow \beta_j \in \partial G', t \rightarrow 1$. Show that $\beta_1 = \beta_2$ if there exists $C \geq 1$ with $k_{G'}(f(x), f(y)) \leq C k_G(x, y)$ for all $x, y \in G$. Show that $\beta_1 = \beta_2$ also holds if there exists $K \geq 1$ such that $M(\Gamma) \leq KM(f\Gamma) \leq K^2M(\Gamma)$ for all curve families Γ in G .

5. Let $D = \{z \in \mathbf{C} : 0 < \arg z < \theta, 0 < |z| < 1\}$, $z_1 = 1, z_2 = e^{i\alpha}$ for $0 < \alpha < \theta, z_3 = e^{i\theta}$ and $z_4 = 0$. Find the modulus of the quadrilateral $(D; z_1, z_2, z_3, z_4)$.

In other words, find a conformal map of D onto $\{z \in \mathbf{C} : \text{Im } z > 0\}$ such that the points z_k are mapped onto the real axis and compute the cross ratio of these points.