

Department of Mathematics and Statistics  
 Riemannian geometry  
 Exercise 1  
 26.1.2016

1. The standard Riemannian metric of the Euclidean space  $\mathbb{R}^n$  is

$$g = \delta_{ij} dx^i dx^j.$$

Write  $g$  in the case of the plane  $\mathbb{R}^2$  by using the polar coordinates  $(r, \vartheta)$ ,  $x^1 = r \cos \vartheta$ ,  $x^2 = r \sin \vartheta$ .

2. A surface of revolution is obtained as follows: Consider a smooth simple (i.e. injective) curve  $\gamma = (r, z): I \rightarrow \mathbb{R}^2$ ,  $\gamma(t) = (r(t), z(t))$ , where  $I \subset \mathbb{R}$  is an interval. By rotating this curve around  $z$ -axis, we get a surface whose cylindrical coordinate representation is

$$(t, \vartheta) \mapsto f(t, \vartheta) = (r(t) \cos \vartheta, r(t) \sin \vartheta, z(t)).$$

We obtain a natural frame  $\partial_t, \partial_\vartheta$  with coframe  $dt, d\vartheta$ . Write the induced Riemannian metric  $dx^2 + dy^2 + dz^2$  from  $\mathbb{R}^3$  in this frame.

3. Let  $(M, g)$  be a Riemannian  $n$ -manifold. Prove that for every  $p \in M$  there exist a neighborhood  $U \ni p$  and smooth vector fields  $E_1, \dots, E_n \in \mathcal{T}(U)$  that form a local orthonormal frame in  $U$ , i.e. the vectors  $(E_1)_q, \dots, (E_n)_q$  form an orthonormal basis for  $T_q M$  for every  $q \in U$ . [Hint: Use the Gram-Schmidt algorithm.]

**Definitions** for problems 4 and 5:

- (a) Let  $(M, g)$ ,  $g = \langle \cdot, \cdot \rangle$ , be a Riemannian manifold,  $U \subset M$  open, and  $f \in C^\infty(U)$  a real-valued function. The *gradient* of  $f$  is a vector field  $\nabla f \in \mathcal{T}(U)$  such that

$$\langle \nabla f, X \rangle = Xf$$

for every  $X \in \mathcal{T}(U)$ .

- (b) We say that a Riemannian metric  $\hat{g} = \langle \langle \cdot, \cdot \rangle \rangle$  is obtained from the Riemannian metric  $g = \langle \cdot, \cdot \rangle$  by a conformal change if

$$(1) \quad \langle \langle \cdot, \cdot \rangle \rangle_p = \varphi(p) \langle \cdot, \cdot \rangle_p$$

for every  $p \in M$ , where  $\varphi: M \rightarrow ]0, \infty[$  is a (positive)  $C^\infty$ -function.

4. Let  $(M, g)$ ,  $g = \langle \cdot, \cdot \rangle$ , be a Riemannian manifold and let  $(U, x)$ ,  $x = (x^1, \dots, x^n)$ , be a chart. Express the gradient,  $\nabla f$ , of a  $C^\infty$ -function  $f: U \rightarrow \mathbb{R}$  in local coordinates, i.e. by using the coordinate frame  $\partial_1, \dots, \partial_n$ .

5. Let  $(M^n, g)$ ,  $g = \langle \cdot, \cdot \rangle$ , be a Riemannian  $n$ -manifold. Suppose that  $\hat{g} = \langle \langle \cdot, \cdot \rangle \rangle$  is obtained from  $g$  by a conformal change (1). Let  $f: M^n \rightarrow \mathbb{R}$  be a  $C^\infty$ -function,  $\nabla f$  its gradient with respect to

the Riemannian metric  $\langle \cdot, \cdot \rangle$ , and  $|\nabla f| = \langle \nabla f, \nabla f \rangle^{1/2}$ . Furthermore, let  $\hat{\nabla} f$  be the gradient of  $f$  with respect to the Riemannian metric  $\langle \langle \cdot, \cdot \rangle \rangle$  and  $\|\hat{\nabla} f\| = \langle \langle \hat{\nabla} f, \hat{\nabla} f \rangle \rangle^{1/2}$ .

- (a) Express  $\hat{\nabla} f$  and  $\|\hat{\nabla} f\|$  in terms of the function  $\varphi$  in formula (1),  $\nabla f$ , and  $|\nabla f|$ .
- (b) Let  $(U, x)$  be a chart. Compare the integrals

$$(2) \quad \int_U |\nabla f|^n d\mu$$

and

$$(3) \quad \int_U \|\hat{\nabla} f\|^n d\hat{\mu}.$$

What do you notice? Above in (2) we integrate with respect to the Riemannian metric  $g$  and in (3) with respect to the Riemannian metric  $\hat{g}$ . In other words,

$$\int_U |\nabla f|^n d\mu = \int_{xU} \left( |\nabla f|^n \sqrt{\det(g_{ij})} \right) \circ x^{-1} dm$$

and

$$\int_U \|\hat{\nabla} f\|^n d\hat{\mu} = \int_{xU} \left( \|\hat{\nabla} f\|^n \sqrt{\det(\hat{g}_{ij})} \right) \circ x^{-1} dm.$$