

Department of Mathematics and Statistics  
Riemannian geometry  
Exercise 8  
5.4.2016

1. Complete the proof of Theorem 6.22: Let  $\gamma : [a, b] \rightarrow M$  be a geodesic. If  $p = \gamma_a$  is not conjugate to  $q = \gamma_b$  and  $v_1 \in T_pM$ ,  $v_2 \in T_qM$ , then there exists a unique Jacobi field  $V$  along  $\gamma$  such that  $V_a = v_1$  and  $V_b = v_2$ . (Uniqueness is proven in the lecture notes.)
2. Let  $M$  be a Riemannian manifold with sectional curvature identically zero. Show that, for every  $p \in M$ , the mapping  $\exp_p : B(0, \varepsilon) \rightarrow B(p, \varepsilon)$  is an isometry whenever  $B(p, \varepsilon)$  is a normal ball at  $p$ .
3. Let  $M$  be complete with  $K(\sigma) \leq 0$  for every 2-planes  $\sigma \subset T_pM$ ,  $\forall p \in M$ . Prove that  $\forall p \in M$ ,  $\exp_p : T_pM \rightarrow M$  is a local diffeomorphism.
4. Let  $M$  be a Riemannian manifold,  $p \in M$ ,  $x, y, z \in T_pM$ ,  $|x| = 1$  and  $\gamma = \gamma^x$ . Let  $Y$  and  $Z$  be Jacobi fields along  $\gamma$  such that  $Y_0 = 0$ ,  $Y'_0 = (D_t Y)_0 = y$ ,  $Z_0 = 0$ , and  $Z'_0 = (D_t Z)_0 = z$ . Prove that

$$\langle Y_t, Z_t \rangle = t^2 \langle y, z \rangle - \frac{t^4}{3} \langle R(y, x)x, z \rangle + O(t^5).$$

5. Let  $e_1, \dots, e_n$  be an orthonormal basis of  $T_pM$ ,  $(U, \varphi)$  the corresponding normal chart at  $p$ , and  $g_{ij}$  the corresponding component functions of the Riemannian metric. Prove that

$$g_{ij}(\exp_p v) = \delta_{ij} - \frac{1}{3} \langle R(e_i, v)v, e_j \rangle + O(|v|^3),$$

for  $\exp_p v \in U$ .