

Department of Mathematics and Statistics  
 Riemannian geometry  
 Exercise 9  
 12.4.2016

1. Let  $M$  be a Riemannian  $n$ -manifold and  $p \in M$ . Show that there exist a neighborhood  $U \subset M$  of  $p$  and smooth vector fields  $E_1, \dots, E_n \in \mathcal{T}(U)$  forming a local orthonormal frame in  $U$  such that, at  $p$ ,  $\nabla_{E_i} E_j(p) = 0$ . [Such a family of vector fields  $E_1, \dots, E_n$  is called a geodesic frame at  $p$ .]

2. Let  $A_{ij} : \mathbb{R} \rightarrow \mathbb{R}$ ,  $i, j = 1, \dots, n$ , be smooth mappings and denote  $A = (A_{ij})$ . Suppose that  $\det A(0) > 0$ . Prove that the function  $\det A$  has the expansion

$$\frac{\det A(t)}{\det A(0)} = 1 + t \cdot \operatorname{tr}(A' A^{-1})(0) + \frac{t^2}{2} \left( \operatorname{tr}(A'' A^{-1})(0) - \operatorname{tr}((A' A^{-1})^2)(0) + (\operatorname{tr}(A' A^{-1})(0))^2 \right) + O(t^3)$$

in a neighborhood of 0.

3. Prove that in the situation of Exercise 8/5,

$$\det(g_{ij}(\exp_p v)) = 1 - \frac{1}{3} \operatorname{Ric}(v, v) + O(|v|^3)$$

for  $\exp_p v \in U$ .

4. Let  $\gamma : I \rightarrow M$  be a geodesic,  $0 \in I$ , and  $p = \gamma(0)$ . Prove that, for every  $h \in C^\infty(p)$ , we have

$$(h \circ \gamma)''(0) = \operatorname{Hess} h(\dot{\gamma}_0, \dot{\gamma}_0).$$

5. Let  $M$  be a Riemannian manifold,  $p \in U \subset M$ , and  $R_0 > 0$  such that  $\exp_p |B(0, R_0) : B(0, R_0) \rightarrow U$  is a diffeomorphism. Define  $\rho : U \rightarrow \mathbb{R}$  by setting  $\rho(x) = d(x, p)$ . Let  $\gamma : [0, R] \rightarrow U$  be a unit speed geodesic,  $\gamma(0) = p$ , and  $0 < R < R_0$ . Let  $r \in ]0, R]$ ,  $X \in T_{\gamma(r)} M$ ,  $|X| = 1$ ,  $\langle X, \dot{\gamma}_r \rangle = 0$ , and let  $\sigma$  be a geodesic such that  $\sigma(0) = \gamma(r)$  and  $\dot{\sigma}(0) = X$ . Furthermore, let  $\Gamma$  be the variation of  $\gamma$ , where  $\Gamma_s$  is the (radial) geodesic from  $p$  to  $\sigma(s)$ . Prove that

$$\operatorname{Hess} \rho(X, X) = \int_0^r (|D_t V|^2 - \langle R(V, \dot{\gamma}) \dot{\gamma}, V \rangle) dt,$$

where  $V$  is the variation field of  $\Gamma$ .