

Exercise 1

Multiplication operators

(These are the central unbounded operators for us: they are used for defining both the free Schrödinger evolution and “interaction potentials”. What is more, every self-adjoint operator A on a separable Hilbert space can be represented, with the aid of spectral theory, as a direct sum of multiplication operators on $L^2(\mu)$, where each μ is an appropriately chosen positive measure on a certain closed subset of \mathbb{R} , the spectrum of A .)

If you do nothing else this week, please try to do this exercise carefully. In case you feel uncertain about working with general measures, you can consider the special case with $X = \mathbb{R}^d$ and μ equal to the Lebesgue measure.)

Let X be a measure space with a positive measure μ , and denote $\mathcal{H} = L^2(\mu)$. For any measurable function $V : X \rightarrow \mathbb{C}$ let M_V denote the *multiplication operator* corresponding to V : define $(M_V\psi)(x) = V(x)\psi(x)$ for $x \in X$ and ψ in

$$D(M_V) := \left\{ \psi \in \mathcal{H} \mid \int_X \mu(dx) |V(x)\psi(x)|^2 < \infty \right\}. \quad (1)$$

Prove that

- (a) M_V is a densely defined operator on \mathcal{H} .
 - (b) M_V is closed. (*Hint*: Recall item (c) on p. 10 of the lecture notes.)
 - (c) $(M_V)^* = M_{V^*}$, where $(V^*)(x) = V(x)^*$, $x \in X$. (Do not forget to check the domains!)
- (*Hint*: For a given $\psi \in \mathcal{H}$, consider $\psi_n(x) = \psi(x)\mathbb{1}_{\{|V(x)| \leq n\}}$ for $n \in \mathbb{N}$.)

Exercise 2

Schrödinger operators do not have bounded extensions

Let $D := C_c^\infty(\mathbb{R}^d)$ be the set of smooth (i.e., arbitrarily many times continuously differentiable) functions with a compact support, which is a dense subspace of $\mathcal{H} = L^2(\mathbb{R}^d)$. Assume $V : \mathbb{R}^d \rightarrow \mathbb{R}$ is smooth and consider the map S defined for $\psi \in D$ by

$$(S\psi)(x) = -\frac{1}{2}\nabla^2\psi(x) + V(x)\psi(x). \quad (2)$$

Then S is an operator with $D(S) = D$ and $R(S) \subset D$.

- (a) Show that S is *symmetric*: $\langle \phi | S\psi \rangle = \langle S\phi | \psi \rangle$ for all $\psi, \phi \in D$.
- (b) Show that, if A extends S , then $A \notin \mathcal{B}(\mathcal{H})$.

(*Hint*: There exists $\psi \in D$ such that $\nabla^2\psi \neq 0$. Consider ψ_λ defined by $\psi_\lambda(x) = \lambda^{\frac{d}{2}}\psi(\lambda x)$, with $\lambda > 0$.)

(Please turn over)

Exercise 3

A weakly continuous unitary semigroup is strongly continuous

Suppose $(U_t)_{t \geq 0}$ is a collection of unitary operators such that $U_0 = 1$, and $U(t+s) = U(t)U(s)$, for all $t, s \geq 0$. Assume, in addition, that $U_t \xrightarrow{w} 1$ when $t \rightarrow 0^+$; that is, assume that $\langle \phi | U_t \psi \rangle \rightarrow \langle \phi | \psi \rangle$ for all $\phi, \psi \in \mathcal{H}$. Show that then also $U_t \psi \rightarrow \psi$ for all $\psi \in \mathcal{H}$, i.e., $U_t \xrightarrow{s} 1$.

Exercise 4

Let S and T be some densely defined, possibly unbounded, operators. Prove the following statements:

- (a) If $S \subset T$, then $T^* \subset S^*$.
- (b) If S is self-adjoint, then S is closed.
- (c) If $S \subset S^*$, then S is symmetric.
- (d) If S is symmetric, then S is closable, its closure $\bar{S} = S^{**}$ is symmetric, and $S \subset S^{**} \subset S^*$.
- (e) Prove that self-adjoint operators are *maximally symmetric*: If S is self-adjoint and T is a symmetric extension of S , then $T = S$.

(*Hint*: You are allowed (and encouraged) to use the results proven in the lecture notes before Theorem 5.10.)

Exercise 5

Let $\Omega \subset \mathbb{R}^d$ and $\mathcal{H} = L^2(\Omega)$ be as in Exercise 1. Assume $V : \Omega \rightarrow \mathbb{R}$ is Lebesgue measurable, and define $u_t : \Omega \rightarrow \mathbb{C}$, $t \geq 0$, by $u_t(x) = \exp(-itV(x))$. Let $U_t = M_{u_t}$.

Show that $(U_t)_{t \geq 0}$ is a strongly continuous unitary semigroup whose infinitesimal generator is M_V . (Hint: Prove first that $|e^{ir} - 1| \leq |r|$ for any $r \in \mathbb{R}$. Exercise 4 might also be helpful.)