

**Stochastic analysis II Exercises-8, 2.5.2017**

1. For  $M \in \mathcal{M}_2^{loc}$  (a local  $L^2$ -martingale) and predictable  $Y(s, \omega)$  such that  $\forall t > 0$

$$\int_0^t Y(s)^2 d\langle M \rangle_s < \infty,$$

let  $X \in \mathcal{M}_2^{loc}$  be the Ito integral process

$$X(t) = (Y \cdot M)_t = \int_0^t Y(s) dM(s)$$

with the property that for every  $N \in \mathcal{M}_2^{loc}$  we have

$$\langle X, N \rangle_t = \int_0^t Y(s) d\langle M, N \rangle$$

Let  $H(s, \omega)$  be bounded predictable process. Since  $X(t) \in \mathcal{M}_2^{loc}$  we can compute the Ito integral  $(H \cdot X)$ .

Use the defining property of the Ito integral to show that

$$I(t) := (H \cdot X)_t = \int_0^t H(s) dX(s) = \int_0^t H(s) Y(s) dM(s).$$

Write the predictable covariation  $\langle I, N \rangle_t$  for  $N \in \mathcal{M}_2^{loc}$ .

2. Use the defining property of the Ito integral to show that the Ito integral is bilinear with respect to the integrand and integrator processes, i.e. if  $M, X \in \mathcal{M}_2^{loc}$ , and  $H, Y$  are bounded predictable processes

$$I(t) = \int_0^t (H(s) + Y(s)) dM(s) = \int_0^t H(s) dM(s) + \int_0^t Y(s) dM(s) =$$

and

$$J(t) = \int_0^t Y(s) d(M + X)(s) = \int_0^t Y(s) dM(s) + \int_0^t Y(s) dX(s)$$

and for  $N \in \mathcal{M}_2^{loc}$  write the predictable covariations  $\langle I, N \rangle_t$  and  $\langle J, N \rangle_t$

3. It is not true that all uniformly integrable local martingales are true martingales. Even local martingales bounded in  $L^2(P)$  need not to be true martingales ! Here we study such counterexample.

Let  $B_t = (B_t^{(1)}, B_t^{(2)}, B_t^{(3)})$  a 3-dimensional brownian motion starting from 0 at time 0, with independent components, so that  $\langle B^{(i)}, B^{(j)} \rangle_t = \delta_{ij}t$ .

The process

$$R_t = |B_t| = \sqrt{\sum_{i=1}^3 (B_t^{(i)})^2}$$

is called the 3-dimensional Bessel process.

- (a) Use Ito formula to compute the semimartingale decomposition of  $R_t$  into a continuous local martingale part  $W_t$  and a continuous process of finite variation.
- (b) Compute  $\langle R \rangle_t = \langle W \rangle_t$  and use Paul Lévy's characterization theorem for Brownian motion to show that the local martingale part of  $R_t$  which satisfies

$$W_t = R_t - \int_0^t \frac{1}{R_s} ds$$

is a Brownian motion in the filtration  $\mathbb{F}$  generated by  $(B_t)$ .

- (c) Show that  $R_t$  is a  $\mathbb{F}$ -submartingale.

Let  $M_t = R_t^{-1}$  for  $t \geq 1$ . We start the process at time 1 since  $R_0 = 0$ .

- (d) Use Ito formula to show that  $(M_t)_{t \geq 1}$  is a local martingale, and write its Ito integral representation.
- (e) Compute  $\langle M \rangle_t$ .
- (f) Show that  $(M_t)$  is a supermartingale. Hint: it is non-negative, you can use Fatou lemma for the sequence of localized martingales  $(M_{t \wedge \tau_n} : t \geq 1)$ ,  $n \in \mathbb{N}$ .
- (g) Let  $\tau_a := \inf\{t \geq 1 : R_t = a\}$ ,  $a > 0$ , with the convention  $\inf\{\emptyset\} = \infty$ .

Show that the stopped process  $(M_t^{\tau_a})_{t \geq 1}$  is a martingale and consequently  $(\tau_{1/n} : n \in \mathbb{N})$

is a localizing sequence for the local martingale  $(M_t : t \geq 1)$

- (h) Let  $0 < r' < y < r''$ . Use the martingale property of  $(M_{t \wedge \tau_r} : t \geq 1)$  to compute  $P(\tau_{r'} < \tau_{r''} | R_1 = y)$ . By the conditioning we mean that we start  $R_t$  at time  $t = 1$  in position  $y$ .
- (i) For  $0 < r < y$  compute also  $P(\tau_r < \infty | R_1 = y)$ .
- (j) Using (g), show that the 3-dimensional Brownian motion is transient,  $|B_t| \rightarrow \infty$   $P$  a.s., meaning that it leaves eventually any ball centered around the origin without coming back, and therefore  $M_\infty = \lim_{t \rightarrow \infty} M_t = 0$ .
- (k) Using the multivariate gaussian density in polar coordinates, compute the probability densities of  $R_t$  and  $M_t$ , and show that the local martingale  $(M_t : t \geq 1)$  is bounded in  $L^2$ , so that in particular it is uniformly integrable. ( We start the martingale at time 1 since there is the square norm explodes near the origin ).

Note however that  $(M_t)_{t \geq 1}$  is not a martingale. Otherwise  $(X_t)$  would be an uniformly integrable martingale so that  $M_t = E(M_\infty | \mathcal{F}_t)$ ,  $t \geq 1$ . But in dimension 3 the Brownian motion is transient, which means that  $M_\infty = 0$ .

- (l) Compute also the probability distribution of  $R_t^2$ .

Show first that for  $t = 1$ ,

$$P(R_1^2 \in dx) = \mathbf{1}(x \geq 0) \frac{1}{\Gamma(3/2)2^{3/2}} \exp(-x/2)x^{\frac{3}{2}-1} dx$$

which is the distribution of a Gamma random variable with shape parameter  $3/2$  and scale parameter  $2$ , (also called chi-square with 3 degrees of freedom and use the scaling property of Brownian motion.

(m) Show that  $E(\langle M \rangle_t) = \infty \forall t \geq 1$ .

**Remark** This is not in contradiction with  $E(M_\infty^2) < \infty$ , since

$$E((M_t - M_1)^2) = E(\langle M \rangle_t - \langle M \rangle_1)$$

holds for martingales but does not need to hold for local martingales. Even if the local martingale  $M_t$  is bounded in  $L^2$ , it means that  $M_t^2$  is bounded in  $L^1$  which does not give the uniform integrability condition which is necessary to take the limit of a localizing sequence under the expectation.

**Remark** In general, when  $M_t$  is a continuous local martingale with localizing sequence  $\tau_n \uparrow \infty$ , to show that it is a true martingale, you need to show that for  $s \leq t$  and  $A \in \mathcal{F}_s$

$$\begin{aligned} E_P(M_{t \wedge \tau_n} \mathbf{1}_A) &\stackrel{?}{\rightarrow} E_P(M_t \mathbf{1}_A) \\ E_P(M_{s \wedge \tau_n} \mathbf{1}_A) &\stackrel{?}{\rightarrow} E_P(M_s \mathbf{1}_A) \end{aligned}$$

where the left sides are equal by since  $(M_{t \wedge \tau_n} : t \geq 0)$  is a martingale. When the local martingale  $(M_t : t \geq 0)$  is uniformly integrable or bounded in  $L^2(P)$ , for fixed  $t$ , the sequence  $(M_{t \wedge \tau_n} : n \in \mathbb{N})$  does not need to be uniformly integrable, which is the condition that we need to take the limit inside the expectation.

Of course if the local martingale is bounded on finite intervals,  $|M_t(\omega)| \leq c(t) < \infty$   $P$  a.s., then it is a true martingale.

Also, when  $M_t \in L^2(P)$  is a martingale and  $\forall t$  and  $Y_t$  is a progressive integrand with

$$E\left(\langle Y \cdot M \rangle\right) = E\left(\int_0^t Y_s^2 d\langle M \rangle_s\right) < \infty \quad \forall t \quad (1)$$

then the Ito integral  $(Y \cdot M)_t \in L^2(P)$  is a true martingale (see Proposition 30 in the lecture notes).

If  $M_t$  is not a square integrable martingale, or  $Y$  does not satisfy the condition 1, the stochastic integral  $(Y \cdot M)_t$  is just local martingale.