

## 10. Potentials for which $H_0 \neq V$ is self-adjoint on $D(H_0)$

Suppose  $A$  and  $B$  are self-adjoint, and  $D(A) \subset D(B)$ . Then  $D(A+B) = D(A) \cap D(B) = D(A)$ , and thus  $A+B$  is densely defined. It is also obviously symmetric. When is  $A+B$  self-adjoint? The following theorem shows that this is the case at least when  $B$  is sufficiently "small" compared to  $A$ .

10.1. Definition: Let  $A$  and  $B$  be

densely defined operators on  $\mathcal{H}$ .  $B$  is said to be  $A$ -bounded, if

a)  $D(B) \supset D(A)$

b)  $\exists \varepsilon, c \in \mathbb{R}$  such that

$$(*) \quad \forall \psi \in D(A): \|B\psi\| \leq \varepsilon \|A\psi\| + c \|\psi\|.$$

If  $B$  is  $A$ -bounded, the infimum  $\varepsilon_0$  of all  $\varepsilon$  for which  $(*)$  holds is called the relative bound of  $B$  with respect to  $A$ , that is,

$$\varepsilon_0 := \inf \{ \varepsilon \in \mathbb{R} \mid \exists c \in \mathbb{R} \text{ s.t. } (*) \text{ holds} \}.$$

\* Typical application would be  $A = H_0$ ,  $B = V$ .

To use above result, it is crucial to check

that  $\forall \psi \in L^2$  for all  $\psi \in D(H_0)$ . If  $V$  is not

bounded, it is important to have some estimates

for generic  $\psi \in D(H_0)$ . For  $\mathcal{H} = L^2(\mathbb{R}^d)$ , ( $d=1$ )

in particular, the functions in  $D(H_0)$  are quite

regular:  $D(-\Delta)$  is classified in Ex. 10.3. and

$D(H_0)$  in Ex. 11.3. Cases  $d \leq 3$  will be discussed in Prop. 10.3.  $\rightarrow$

### 10.2. Theorem (Kato-Rellich)

Let  $A$  be self-adjoint,  $S$  densely defined and symmetric, and suppose  $S$  is  $A$ -bounded with a relative bound  $\epsilon_0 < 1$ .

Then  $A+S$  is self-adjoint on  $D(A)$ , and any core  $D^2$  of  $A$  is also a core of  $A+S$ .

Proof: Since  $D(S) \supset D(A)$ , now  $D(A+S) = D(A)$  is the natural domain of  $A+S$ . We will prove that  $\exists \mu_0 > 0$  s.t.  $R(A+S \pm i\mu_0) = \mathcal{H}$ .  
 $\Rightarrow R(\frac{1}{\mu_0}(A+S) \pm i) = \mathcal{H}$ . Let  $T = \frac{1}{\mu_0}(A+S)$ .

Then  $D(T) = D(A) \subset D(S) \Rightarrow \forall \phi, \psi \in D(T)$ :  
 $(\phi, T\psi) = \frac{1}{\mu_0} (\phi, A\psi + S\psi)$   
 $= \frac{1}{\mu_0} [(\phi, A\psi) + (\phi, S\psi)]$   
 $\stackrel{A, S \text{ symm.}}{=} \frac{1}{\mu_0} (A\phi + S\phi, \psi) = (T\phi, \psi)$ .

Thus  $T$  is densely defined, and symmetric, with  $R(T \pm i) = \mathcal{H} \Rightarrow C_T$  is unitary.

Thus by Lemma 9.10.  $T$  is self-adjoint.  
 $\Rightarrow A+S = \mu_0 T$  is self-adjoint. (on  $D(A)$ ).

[In general, if  $\lambda \in \mathbb{C}$ , and  $T$  is densely def., then  $D(\lambda T) = D(T)$  and  $(\lambda T)^* = \lambda^* T^*$ .  
Proof: left as exercise.]

Thus to prove the first statement, we only need to find  $\mu_0 > 0$  for which  $R(A+S \pm i\mu_0) = \mathcal{H}$ .

For any  $\mu \in \mathbb{R}, \mu \neq 0$ , since  $A$  is self-adjoint,  
(\*)  $\|(A+i\mu)\psi\|^2 = \|A\psi\|^2 + \mu^2 \|\psi\|^2 \quad \forall \psi \in D(A)$ .  
 $\Rightarrow A+i\mu$  is 1-1  $\Rightarrow \exists T := (A+i\mu)^{-1} \in R(A+i\mu) \rightarrow D(A)$ .  
 $\Rightarrow AT = R(A+i\mu) \rightarrow R(A)$  and we have  
by (\*),  $\forall \phi \in D(T) : \|\phi\|^2 = \|(A+i\mu)T\phi\|^2$   
 $= \|AT\phi\|^2 + \mu^2 \|T\phi\|^2$ .

(10)

$$\Rightarrow \|AT\phi\| \leq \|\phi\|, \quad \|T\phi\| \leq \frac{1}{|\mu|} \|\phi\| \quad \forall \phi \in D(T).$$

$$\Rightarrow \|AT\| \leq 1, \quad \|T\| \leq \frac{1}{|\mu|}.$$

By assumption,  $\exists 0 \leq \varepsilon < 1$  and  $c \in \mathbb{R}$  s.t.

$$\forall \psi \in D(A) : \|S\psi\| \leq \varepsilon \|A\psi\| + c \|\psi\|$$

$\psi = T\phi$

$$\Rightarrow \forall \phi \in D(T) : \|ST\phi\| \leq \varepsilon \|AT\phi\| + c \|T\phi\|$$

$$\leq \left(\varepsilon + \frac{c}{|\mu|}\right) \|\phi\|.$$

Thus, if we set  $\mu$  equal to  $\mu_0 > \max\left(0, \frac{c}{1-\varepsilon}\right)$ ,

we have  $\mu_0 > 0$  and  $a = \varepsilon + \frac{c}{\mu_0} < 1$ .

Let  $T_0 = (A + i\mu_0)^{-1}$ , when we have proven that  $\|ST_0\phi\| \leq a\|\phi\|$ , with  $0 \leq a < 1$ .

9.10.  $\Rightarrow$  Since  $A$  is self-adjoint,  $\frac{1}{\mu_0}A$  is also s.a.,  
 $\mathcal{R} = \mathcal{R}\left(\frac{1}{\mu_0}A + i\right) = \mathcal{R}(A + i\mu_0) = D(T_0)$ .

Thus  $T_0 : \mathcal{R} \rightarrow D(A)$  and  $D(ST_0) = \mathcal{R}$ .

Therefore,  $ST_0 \in \mathcal{B}(\mathcal{R})$  with  $\|ST_0\| \leq a < 1$ .

Let  $B = \sum_{n=0}^{\infty} (-ST_0)^n$ , which by  $\|ST_0\| < 1$

is a norm-convergent sum in  $\mathcal{B}(\mathcal{R})$

$\Rightarrow B \in \mathcal{B}(\mathcal{R})$ . For any  $\phi \in \mathcal{R}$ , it is obvious that  $ST_0 B\phi = -\sum_{n=1}^{\infty} (-ST_0)^n \phi$   
 $= -(B\phi - \phi)$

$\Rightarrow \phi = -(1 + ST_0)B\phi$ . (In fact,  $B = (1 + ST_0)^{-1}$ .)

Thus  $\mathcal{R}(1 + ST_0) = \mathcal{R}$ .

On the other hand,  $\forall \psi \in D(A)$ ,

$$(1 + ST_0)(A + i\mu_0)\psi$$

$$= (A + i\mu_0)\psi + S\psi = (A + S + i\mu_0)\psi.$$

Since  $\mathcal{R}(A + i\mu_0) = \mathcal{R} \Rightarrow \mathcal{R}(A + S + i\mu_0)$   
 $= \mathcal{R}(1 + ST_0) = \mathcal{R}$ .

The proof, that also  $\mathcal{R}(A + S - i\mu_0) = \mathcal{R}$ , is essentially identical. (Consider  $T'_0 = (A - i\mu_0)^{-1}$  and use  $\|(A - i\mu_0)\psi\|^2 = \|A\psi\|^2 + \mu_0^2 \|\psi\|^2$

and  $\mathcal{R}(A - i\mu_0) = \mathcal{R}$  to prove  $\|ST'_0\| \leq a < 1$ ,

$\mathcal{R}(1 + ST'_0) = \mathcal{R} \Rightarrow \mathcal{R}(A + S - i\mu_0) = \mathcal{R}$ .)

Therefore, we have proven that  $A + S$  is self-adjoint.

Suppose then that  $D'$  is a core of  $A$ .  
Then  $S' := (A+S)|_{D'}$  is densely defined  
and symmetric.  $\Rightarrow \overline{S'} \subset A+S$ .

Suppose  $\psi \in D(A+S) \Rightarrow \psi \in D(A) \subset D(S)$ .

Since  $A$  is closed, and  $D'$  is its core,

$\exists$  sequence  $\psi_n \in D'$  s.t.  $\psi_n \rightarrow \psi$  and

$A\psi_n \rightarrow A\psi$ . Then  $\forall n: \psi - \psi_n \in D(S)$  and

$$\|S(\psi - \psi_n)\| \leq \epsilon \|A\psi - A\psi_n\| + c \|\psi - \psi_n\| \rightarrow 0, \text{ when } n \rightarrow \infty.$$

and thus  $S\psi_n \rightarrow S\psi$ .

Since  $S'\psi_n = A\psi_n + S\psi_n, \forall n$ ,

we have also  $\overline{S'}\psi_n \rightarrow A\psi + S\psi$ .

Since  $\overline{S'}$  is closed, this implies (by (cc))

$\psi \in D(\overline{S'})$  and  $\overline{S'}\psi = A\psi + S\psi = (A+S)\psi$ .

Therefore,  $A+S \subset \overline{S'}$  and  $\overline{S'} \subset A+S$

$\Rightarrow \overline{S'} = A+S$  and we have proven that  
 $D'$  is a core of  $A+S$ .  $\square$

Remark: For any  $A, S$  as in Thm. 10.2.  
( $A$  self-adj. &  $S$  symm.)

It is sufficient to check that (\*)  
holds on some core of  $A$ . That is,  
if  $D'$  is a core of  $A, D(A) \subset D(S)$

and  $\exists \epsilon, c \in \mathbb{R}$  s.t.  $\forall \psi \in D'$

$$(**) \quad \|S\psi\| \leq \epsilon \|A\psi\| + c \|\psi\|,$$

then (\*\*) holds for all  $\psi \in D(A)$ .

To see this, note that since  $A$  is closed,

for any  $\psi \in D(A) \Rightarrow \exists \psi_n \in D', \psi_n \rightarrow \psi, A\psi_n \rightarrow A\psi$ .

Thus  $\|S\psi_n\| \leq \epsilon \|A\psi_n\| + c \|\psi_n\| \forall n$

$$\xrightarrow{n \rightarrow \infty} \epsilon \|A\psi\| + c \|\psi\|$$

Also  $\forall \phi \in D(S): (\phi, S\psi_n) = (S\phi, \psi_n)$

$$\xrightarrow{n \rightarrow \infty} (\phi, S\psi) = (\phi, S\psi) \text{ since } \psi \in D(S).$$

Since  $D(S)$  is dense and  $\|S\psi_n\|$  is bounded

$\Rightarrow \forall \phi \in \mathcal{H}: (\phi, S\psi_n) \rightarrow (\phi, S\psi)$ . [exercise]

$$\Rightarrow (S\psi, S\psi_n) \rightarrow \|S\psi\|^2$$

$$\begin{aligned} \Rightarrow 0 \leq \|S\psi - S\psi_n\|^2 &= \|S\psi\|^2 + \|S\psi_n\|^2 - 2\operatorname{Re}(S\psi, S\psi_n) \\ &\leq \|S\psi\|^2 - 2\operatorname{Re}(S\psi, S\psi_n) + (\epsilon \|A\psi_n\| + c \|\psi_n\|)^2 \\ &\xrightarrow{n \rightarrow \infty} \|S\psi\|^2 - 2\|S\psi\|^2 + (\epsilon \|A\psi\| + c \|\psi\|)^2 \end{aligned}$$

$$\Rightarrow \|S\psi\| \leq \epsilon \|A\psi\| + c \|\psi\| \quad \square$$

10.3. Proposition Suppose  $d \leq 3$  and

$\mathcal{H} = L^2(\mathbb{R}^d)$ . Then for all  $\epsilon > 0$  there is  $C_\epsilon > 0$  s.t.

$$\|u\|_\infty \leq \epsilon \|H_0 u\| + C_\epsilon \|u\| \quad \forall u \in D(H_0)$$

where  $H_0 = -\frac{1}{2}\nabla^2$  = free Hamiltonian.

Proof Exercise.  $\square$

The following theorem now covers most one-particle potentials which do not grow at infinity.

10.4. Theorem Suppose  $d \leq 3$ , and

$V_1 \in L^\infty(\mathbb{R}^d)$ ,  $V_2 \in L^2(\mathbb{R}^d)$  are such

that  $V = V_1 + V_2$  is real-valued.

then  $H = H_0 + V = -\frac{1}{2}\nabla^2 + M_V$  is self-adjoint on  $D(-\nabla^2)$

and essentially self-adjoint on  $S$ , (and on  $\mathcal{D}$ )

Proof. Since  $V$  is real,  $M_V$  is self-adjoint with  $D(M_V) = \{Vu \in L^2\}$ .

If  $u \in L^2$  is such that  $\|u\|_\infty < \infty$  then

$$\begin{aligned} \int dx |V_1(x)u(x)|^2 &\leq \|V_1\|_\infty^2 \|u\|^2, \\ \int dx |V_2(x)u(x)|^2 &\leq \|u\|_\infty^2 \|V_2\|_{L^2}^2. \\ \Rightarrow Vu = V_1u + V_2u &\in L^2 \text{ and} \\ \|Vu\| &\leq \|V_1\|_\infty \|u\| + \|u\|_\infty \|V_2\|. \end{aligned}$$

By proposition 10.3, if  $u \in D(H_0)$ , then  $\|u\|_\infty < \infty$ . Therefore  $D(H_0) \subset D(M_V)$ .

In addition,  $\forall \epsilon > 0 \exists C_\epsilon \geq 0$  s.t.  $\forall u \in D(H_0)$

$$\begin{aligned} \|M_V u\| &\leq \|V_1\|_\infty \|u\| \\ &\quad + \|V_2\|_{L^2} (\epsilon \|H_0 u\| + C_\epsilon \|u\|) \\ &= \|V_2\|_{L^2} \epsilon \|H_0 u\| + (\|V_1\|_\infty + C_\epsilon \|V_2\|_{L^2}) \|u\|. \end{aligned}$$

This proves that  $M_V$  is  $H_0$ -bounded with relative bound 0. In particular, since  $H_0$  and  $M_V$  are self-adjoint, we can apply the Kato-Rellich Theorem,

$\Rightarrow H_0 + M_V$  is self-adjoint on  $D(H_0) = D(-\Delta) = D(-\nabla^2)$ . Since  $S_d$  is a core of  $-\nabla^2$  (Proposition 6.6.4.),  $H_0 + M_V$  is essentially self-adj. on  $S$ .  $\square$

10.5. Theorem: Suppose  $V \in L^1_{loc}(\mathbb{R}^d)$

and there is  $M \in \mathbb{R}$  s.t.  $V(x) \geq -M$  for Lebesgue a.r.  $x \in \mathbb{R}^d$ . Let  $D(H)$  collect all  $\psi \in L^2(\mathbb{R}^d)$  for which  $\forall \psi \in L^1_{loc}(\mathbb{R}^d)$  and  $\exists \varphi_\psi \in L^2(\mathbb{R}^d)$  s.t.

$$(f, \varphi_\psi) = (-\frac{1}{2}\Delta f, \psi) + \int dx f(x)^* V(x)\psi(x) \quad \forall f \in \mathcal{D}(\mathbb{R}^d).$$

Then  $\varphi_\psi$  is unique, and setting  $H\psi := \varphi_\psi$  for  $\psi \in D(H)$  defines a self-adjoint operator on  $L^2(\mathbb{R}^d)$ .

Proof: Uses quadratic forms, see Reed & Simon: book II, Theorem X.32.  $\square$

\* For instance, if  $V \in C(\mathbb{R}^d)$  and  $\inf V \geq -M$ , then  $\mathcal{D}_d \subset D(H)$  but it can happen that  $S_d \not\subset D(H)$ : if  $\psi \in S_d$  and  $\varphi_\psi(x) := -\frac{1}{2}\Delta\psi(x) + V(x)\psi(x)$ , then  $V\psi \in L^1_{loc}$  and for any  $f \in \mathcal{D}$  we have  $f^* \varphi_\psi \in L^1$

$$\text{with } \int dx f(x)^* \varphi_\psi(x) = (f, -\frac{1}{2}\Delta\psi) + \int dx f^* V\psi \\ = (-\frac{1}{2}\Delta f, \psi) + \int dx f^* V\psi.$$

Thus if  $\psi \in \mathcal{D}_d$  also  $\int dx |V\psi|^2 \leq \max_{x \in K} |V(x)\psi(x)|^2 \cdot |K| < \infty$   
 $K = \text{supp } \psi$ .

$\Rightarrow \varphi_\psi \in L^2$  and hence  $\psi \in D(H)$ .

But if  $V$  increases too fast at  $\infty$ , for instance, exponentially, then  $\exists \psi \in S_d$  s.t.  $\varphi_\psi \notin L^2$ .