

**Introduction to mathematical physics:  
Quantum dynamics**

Homework set 3  
Monday 23.9.2013

**Exercise 1**

Every  $\phi, \psi \in \mathcal{H}$  satisfies a *polarization identity*:

$$(\phi, \psi) = \frac{1}{4} (\|\phi + \psi\|^2 - \|\phi - \psi\|^2 - i\|\phi + i\psi\|^2 + i\|\phi - i\psi\|^2) .$$

The identity can be generalized to include an action with a bounded operator: for all  $\phi, \psi \in \mathcal{H}$  and  $A \in \mathcal{B}(\mathcal{H})$

$$(\phi, A\psi) = \frac{1}{4} \sum_{n=0}^3 i^{-n} (\phi + i^n \psi, A(\phi + i^n \psi)) . \quad (1)$$

(The polarization identity clearly corresponds to the case  $A = 1$ .)

- (a) Prove the *generalized polarization identity* in (1).
- (b) Suppose  $T \in \mathcal{B}(\mathcal{H})$ . Show that, if  $(\psi, T\psi) = 0$  for all  $\psi \in \mathcal{H}$ , then  $T = 0$ . (This implies that the “expectation values” of  $T$  determine  $T$  uniquely.)
- (c) An operator  $T : D \rightarrow \mathcal{H}$  is called *positive* if  $(\psi, T\psi) \geq 0$  for all  $\psi \in D$ . Prove that, if  $T \in \mathcal{B}(\mathcal{H})$  is positive, then it is self-adjoint.

**Exercise 2**

Let  $M \subset \mathcal{H}$  be a subspace. Show that  $(M^\perp)^\perp = \overline{M}$ .

**Exercise 3**

Let  $T, S \in \mathcal{B}(\mathcal{H})$ , and  $\alpha \in \mathbb{C}$  be arbitrary. Show that all of the following statements hold for the related adjoint operators.

- (a)  $(T + S)^* = T^* + S^*$
- (b)  $(\alpha T)^* = \alpha^* T^*$
- (c)  $(ST)^* = T^* S^*$  (notation:  $ST = S \circ T$ )
- (d)  $T^{**} = T$  (notation:  $T^{**} = (T^*)^*$ )
- (e)  $\|T^* T\| = \|T\|^2$

This proves that “ $*$ ” is an involution on  $\mathcal{B}(\mathcal{H})$  which makes it into a  $C^*$ -algebra.

**Exercise 4**

Suppose  $U \in \mathcal{B}(\mathcal{H})$ . Show that the following statements are equivalent:

- (a)  $U$  is a unitary operator:  $U^* U = 1 = U U^*$ .
- (b)  $R(U) = \mathcal{H}$  and  $(U\psi, U\phi) = (\psi, \phi)$  for all  $\psi, \phi \in \mathcal{H}$ .
- (c)  $R(U) = \mathcal{H}$  and  $\|U\psi\| = \|\psi\|$  for all  $\psi \in \mathcal{H}$ .

(Please turn over)

## Exercise 5

Two Hilbert spaces  $\mathcal{H}_1, \mathcal{H}_2$  are said to be isomorphic, if there exists a *unitary map* between them (a map is unitary if it is linear, invertible, and preserves the scalar product). We denote this by  $\mathcal{H}_1 \cong \mathcal{H}_2$ . Show that

- (a)  $L^2(\mathbb{R}^3) \otimes \mathbb{C}^2 \cong L^2(\mathbb{R}^3, \mathbb{C}^2) \cong L^2(\mathbb{R}^3) \oplus L^2(\mathbb{R}^3)$ . (Here  $L^2(\mathbb{R}^3, \mathbb{C}^2)$  denotes the  $L^2$ -space of *two-component wavefunctions*,  $\psi : \mathbb{R}^3 \rightarrow \mathbb{C}^2$ , with a scalar product  $(\phi, \psi) = \int dx \sum_{i=1}^2 \phi_i(x)^* \psi_i(x)$ .)
- (b)  $L^2([0, 1]^d) \otimes L^2([0, 1]^{d'}) \cong L^2([0, 1]^{d+d'})$  for any  $d, d' \in \mathbb{N}_+$ . (Hint: Fourier series.)