

University of Helsinki / Department of Mathematics and Statistics  
 SCIENTIFIC COMPUTING  
 Exercise 04, 29.9.2014

Problem sessions will be held on Monday at 16-18, B322.

N.B. The files mentioned in the exercises (if any) are available on the course homepage

1. (a) Let  $X$  and  $Y$  be independent uniformly distributed random variables on  $(0, 1)$ . As we know, samples of  $X$  can be generated by  $x = \text{rand}(1, 100)$ ; for instance. Now it is a basic fact (this need not be proven) that the new random variables

$$U = \cos(2\pi X)\sqrt{-2\log Y}; \quad V = \sin(2\pi X)\sqrt{-2\log Y}$$

follow the normal distribution with parameters  $(0, 1)$ , i.e. with mean 0 and variance 1. Use this so called Box-Müller method to generate 200 samples of normal distribution, plot the result with the command `hist`, compute the mean and standard deviation of the sample.

(b) The amplitude distribution of a signal sent by a mobile phone to a base station follows so called Rayleigh distribution. Suppose that  $X_1, X_2$  are zero-mean normally distributed random variables with variance  $\sigma^2$  and define a new random variable  $R$  by  $R = \sqrt{X_1^2 + X_2^2}$ . Then  $R$  follows the Rayleigh distribution. Generate 100 samples of a Rayleigh distribution and plot the histogram.

2. Suppose that  $f: [a, b] \rightarrow [0, \infty)$  is continuous and that  $0 \leq f(x) \leq M$  for all  $x \in [a, b]$ . Use the Monte Carlo method to approximate the value of

$$\int_a^b f(x) dx,$$

that is, choose  $m$  random points in  $[a, b] \times [0, M]$  and compute the ratio  $p/m$  where  $p$  is the number of points below the graph of  $f(x)$ . Apply this method for the function

$$f(x) = \sum_{j=1}^n c_j(1 + \sin(d_j x))$$

FILE: ~/mme11/demo11/d04/d04.tex — 21. elokuuta 2014 (klo 11.35).

in  $[0, 1]$  with  $m = 10j$ ,  $j = 10 : 10 : 100$  where  $n = 4$ ,  $c = \text{rand}(1, n)$ ,  $d = 1 + 3 * \text{rand}(1, n)$ . Compare your result to the exact value

$$\int_a^b f(x) dx = (b - a) \sum_{j=1}^n c_j + \sum_{j=1}^n (c_j / d_j) (\cos(d_j * a) - \cos(d_j * b)),$$

see Problem 3/Exercise 2.

3. The ASCII codes of capital letters A,...,Z are 65,...,90. A simple ciphering method, so called Caesar cipher, is the following. Fix an integer  $p \in [1, 25]$ . Each letter is replaced by another, obtained by increasing its ASCII code by the constant  $p$ . (Note that we recycle: 91 corresponds to 65 i.e. after Z come A,B,C,...). The program `hlp043.m` shows how this happens. Use this idea to decipher the message:

Q C A D I H C S F U C G I A

4. We want to fit a model of the form  $f(x) = ae^{bx}$  to the data set

x	1	3	4	6	9	15
y	4.0	3.5	2.9	2.5	2.75	2.0

where  $a$  and  $b$  are parameters to be determined from the data.

(a) For this purpose we introduce new transformed variables  $X = x$ ,  $Y = \log(y)$ . Carry out this data transformation and print out the transformed variables.

(b) After the transformation the new model is  $F(x) = \log f(x) = bx + \log a$ . Apply the usual LSQ method to find  $b$  and  $\log a$ .

(c) Print the results in the following format

x(i)	y(i)	Y(i)	a*exp(b*x(i))	y(i)-a*exp(b*x(i))
1	4.0	....		
.....				
15	2.0			

and plot the data and the fitted curve in the same figure.

5. For a complex  $n \times n$  matrix  $a$  let  $P_i = \sum_{j=1, j \neq i}^n |a_{i,j}|$ ,  $m_0 = \min\{|a_{i,i}| - P_i : i = 1, \dots, n\}$ ,  $m = \max\{m_0, 0\}$ ,  $M = \max\{|a_{i,i}| + P_i : i = 1, \dots, n\}$ .

By Gerschgorin's theorem (recall Exercise 03) the eigenvalues  $\lambda_i$  of  $a$  satisfy

$$m \leq |\lambda_i| \leq M; \quad i = 1, \dots, n$$

and it also follows that  $m^n \leq D \leq M^n$ ,  $D = |\det(a)|$ . Set  $m_1 = \min\{|\lambda_i| : i = 1, \dots, n\}$  and  $m_2 = \max\{|\lambda_i| : i = 1, \dots, n\}$ . Write a MATLAB script that experimentally confirms these statements, by printing out the test results in the following format

```
n   m   m1   m2   M   D - m^n   M^n -D
```

Use random complex  $n \times n$  matrices,  $n=5:5:50$ .

Repeat the experiment for the matrices  $a=2*n*eye(n)+rand(n,n)+i*rand(n,n)$ .

6. The arithmetic-geometric mean  $ag(a, b)$  of two positive numbers  $a > b > 0$  is defined as  $ag(a, b) = \lim a_n$ , where  $a_0 = a$ ,  $b_0 = b$ , and

$$a_{n+1} = (a_n + b_n)/2, \quad b_{n+1} = \sqrt{a_n b_n}, \quad n = 0, 1, 2, \dots$$

(a) Write a function, which takes two arguments (double), computes  $ag$  and returns the value (double).

(b) The hypergeometric function  ${}_2F_1(a, b; c; x)$  is defined as a sum of the series,

$$\begin{aligned} {}_2F_1(a, b; c; x) = & 1 + \frac{abx}{c} \frac{1}{1!} + \frac{a(a+1)b(b+1)x^2}{c(c+1)} \frac{1}{2!} + \dots \\ & + \frac{a(a+1)\dots(a+j-1)b(b+1)\dots(b+j-1)x^j}{c(c+1)\dots(c+j-1)} \frac{1}{j!} + \dots \end{aligned}$$

This hypergeometric series converges for  $\text{abs } x < 1$ . Gauss proved in 1799 that there is a connection between the hypergeometric function and the arithmetic-geometric mean,

$${}_2F_1\left(\frac{1}{2}, \frac{1}{2}; 1; r^2\right) = \frac{1}{ag(1, \sqrt{1-r^2})}$$

for  $0 < r < 1$ . Tabulate the difference of the two sides of this identity for  $r = 0.05k$ ,  $k = 1, \dots, 19$ . Use the routine on the web-page to calculate the values of the  ${}_2F_1$  or the MATLAB built-in program (`help hypergeom`).