

Building a matrix model for one-dimensional convolution

Samuli Siltanen

Department of Mathematics and Statistics
University of Helsinki, Finland
samuli.siltanen@helsinki.fi
www.siltanen-research.net

Applications of Matrix Computations

University of Helsinki, Finland

October 5, 2016

Outline

The convolution process

Constructing the matrix

We define discrete convolution using periodic boundary conditions

Let $p \in \mathbb{R}^n$ and $s \in \mathbb{R}^n$. We call s the *signal* and p the *point spread function (PSF)*.


Convolution, denoted by $p * s \in \mathbb{R}^n$, is defined by the formula

$$(p * s)_j = \sum_{\ell=1}^n p_{\ell} s_{j-\ell}, \quad (1)$$


where $s_{j-\ell}$ is defined by periodic extension for the cases $j - \ell < 1$ and $j - \ell > n$. For example, $s_0 = s_n$ and $s_{-1} = s_{n-1}$ and $s_{n+1} = s_1$.

Step-by-step construction of the convolution vector (1) is shown in the following slides for a special case with $n = 10$. Note that only nonzero elements of $p \in \mathbb{R}^n$ are plotted. The elements $(p * s)_j$ are shown in blue color; the last slide (position 10) shows the complete vector $p * s \in \mathbb{R}^n$.


Convolution, position 1




Convolution, position 2




Convolution, position 3




Convolution, position 4




Convolution, position 5




Convolution, position 6




Convolution, position 7




Convolution, position 8



Convolution, position 9



Convolution, position 10



Outline

The convolution process

Constructing the matrix


Given a PSF vector p , the mapping $\mathcal{A} : \mathbb{R}^n \rightarrow \mathbb{R}^n$ defined by $\mathcal{A}(s) = p * s$ is linear

Therefore, we can construct a $n \times n$ matrix A so that


$$\mathcal{A}(s) = As.$$

The following slides demonstrate the construction of the matrix A . Note that the structure of A is highly redundant as the same PSF elements appear many times.


The Matlab command `convmtx.m` is useful for building convolution matrices.




0.50 0.10 0.05 0 0 0 0 0 0.05 0.30




0.50	0.10	0.05	0	0	0	0	0	0.05	0.30
0.30	0.50	0.10	0.05	0	0	0	0	0	0.05




0.50	0.10	0.05	0	0	0	0	0	0.05	0.30
0.30	0.50	0.10	0.05	0	0	0	0	0	0.05
0.05	0.30	0.50	0.10	0.05	0	0	0	0	0




0.50	0.10	0.05	0	0	0	0	0	0.05	0.30
0.30	0.50	0.10	0.05	0	0	0	0	0	0.05
0.05	0.30	0.50	0.10	0.05	0	0	0	0	0
0	0.05	0.30	0.50	0.10	0.05	0	0	0	0




0.50	0.10	0.05	0	0	0	0	0	0.05	0.30
0.30	0.50	0.10	0.05	0	0	0	0	0	0.05
0.05	0.30	0.50	0.10	0.05	0	0	0	0	0
0	0.05	0.30	0.50	0.10	0.05	0	0	0	0
0	0	0.05	0.30	0.50	0.10	0.05	0	0	0




0.50	0.10	0.05	0	0	0	0	0	0.05	0.30
0.30	0.50	0.10	0.05	0	0	0	0	0	0.05
0.05	0.30	0.50	0.10	0.05	0	0	0	0	0
0	0.05	0.30	0.50	0.10	0.05	0	0	0	0
0	0	0.05	0.30	0.50	0.10	0.05	0	0	0
0	0	0	0.05	0.30	0.50	0.10	0.05	0	0




0.50	0.10	0.05	0	0	0	0	0	0.05	0.30
0.30	0.50	0.10	0.05	0	0	0	0	0	0.05
0.05	0.30	0.50	0.10	0.05	0	0	0	0	0
0	0.05	0.30	0.50	0.10	0.05	0	0	0	0
0	0	0.05	0.30	0.50	0.10	0.05	0	0	0
0	0	0	0.05	0.30	0.50	0.10	0.05	0	0
0	0	0	0	0.05	0.30	0.50	0.10	0.05	0



0.50	0.10	0.05	0	0	0	0	0	0.05	0.30
0.30	0.50	0.10	0.05	0	0	0	0	0	0.05
0.05	0.30	0.50	0.10	0.05	0	0	0	0	0
0	0.05	0.30	0.50	0.10	0.05	0	0	0	0
0	0	0.05	0.30	0.50	0.10	0.05	0	0	0
0	0	0	0.05	0.30	0.50	0.10	0.05	0	0
0	0	0	0	0.05	0.30	0.50	0.10	0.05	0
0	0	0	0	0	0.05	0.30	0.50	0.10	0.05



0.50	0.10	0.05	0	0	0	0	0	0.05	0.30
0.30	0.50	0.10	0.05	0	0	0	0	0	0.05
0.05	0.30	0.50	0.10	0.05	0	0	0	0	0
0	0.05	0.30	0.50	0.10	0.05	0	0	0	0
0	0	0.05	0.30	0.50	0.10	0.05	0	0	0
0	0	0	0.05	0.30	0.50	0.10	0.05	0	0
0	0	0	0	0.05	0.30	0.50	0.10	0.05	0
0	0	0	0	0	0.05	0.30	0.50	0.10	0.05
0.05	0	0	0	0	0	0.05	0.30	0.50	0.10



0.50	0.10	0.05	0	0	0	0	0	0.05	0.30
0.30	0.50	0.10	0.05	0	0	0	0	0	0.05
0.05	0.30	0.50	0.10	0.05	0	0	0	0	0
0	0.05	0.30	0.50	0.10	0.05	0	0	0	0
0	0	0.05	0.30	0.50	0.10	0.05	0	0	0
0	0	0	0.05	0.30	0.50	0.10	0.05	0	0
0	0	0	0	0.05	0.30	0.50	0.10	0.05	0
0	0	0	0	0	0.05	0.30	0.50	0.10	0.05
0.05	0	0	0	0	0	0.05	0.30	0.50	0.10
0.10	0.05	0	0	0	0	0	0.05	0.30	0.50