

DEFORMATION QUANTIZATION AND BOREL'S THEOREM IN LOCALLY CONVEX SPACES

MIROSLAV ENGLIŠ, JARI TASKINEN

ABSTRACT. It is well known that one can often construct a star-product by expanding the product of two Toeplitz operators asymptotically into a series of another Toeplitz operators multiplied by increasing powers of the Planck constant h . This is the Berezin-Toeplitz quantization. We show that one can obtain in a similar way in fact any star-product which is equivalent to the Berezin-Toeplitz star-product, by using instead of Toeplitz operators suitable other assignments from compactly supported smooth functions to bounded linear operators on the corresponding Hilbert spaces. A crucial ingredient in the proof is the generalization, due to Colombeau, of the classical theorem of Borel on the existence of a function with prescribed derivatives of all orders at a point, which reduces the proof to a construction of a locally convex space enjoying some special properties.

1. INTRODUCTION AND BACKGROUND

Let Ω be a domain in $\mathbf{C}^n \cong \mathbf{R}^{2n}$ equipped with a Poisson bracket $\{\cdot, \cdot\}$, i.e. a first order bidifferential operator

$$(1) \quad \{f, g\}(x) = \sum_{j,k=1}^{2n} B_{jk}(x) \frac{\partial f}{\partial x_j} \frac{\partial g}{\partial x_k}, \quad x \in \mathbf{R}^{2n},$$

where, for each $x \in \Omega$, the matrix $B_{jk}(x)$ is skew-symmetric and nondegenerate. Denote by $C^\infty(\Omega)[[h]]$ the ring of formal power series in a variable h with coefficients in $C^\infty(\Omega)$. A *star product* on Ω is a $\mathbf{C}[[h]]$ -bilinear map $*$: $C^\infty(\Omega)[[h]] \times C^\infty(\Omega)[[h]] \rightarrow C^\infty(\Omega)[[h]]$ such that

- (i) $*$ is associative,
- (ii) there exist bidifferential operators $C_j : C^\infty(\Omega) \times C^\infty(\Omega) \rightarrow C^\infty(\Omega)$ such that

$$(2) \quad f * g = \sum_{j=0}^{\infty} h^j C_j(f, g), \quad \forall f, g \in C^\infty(\Omega);$$

- (iii) the operators C_j satisfy

$$\begin{aligned} C_0(f, g) &= fg, \\ C_1(f, g) - C_1(g, f) &= \frac{i}{2\pi} \{f, g\}, \end{aligned}$$

and

$$C_j(f, \mathbf{1}) = C_j(\mathbf{1}, f) = 0, \quad \forall j \geq 1.$$

2000 *Mathematics Subject Classification.* Primary 46A13; Secondary 53D55, 26B05, 47B35.

Key words and phrases. Berezin-Toeplitz quantization, Borel theorem, Frechet space, inductive limit.

Research of the first author supported by GA ĀR grant no. 201/03/0041 and by AV ĀR IRP no. AV0Z10190503.

Note that the last equality means precisely that $\mathbf{1}$ is the unit element for $*$.

Remark. Generalization to complex manifolds Ω is straightforward. \square

Two star products $*, *'$ are called *equivalent* if there exists a sequence of linear differential operators M_0, M_1, M_2, \dots on $C^\infty(\Omega)$ with $M_0 = I$ (the identity operator) such that we have the following equality of formal power series

$$(3) \quad M(f *' g) = Mf * Mg, \quad \forall f, g \in C^\infty(\Omega)[[h]],$$

where

$$(4) \quad Mf = \sum_{j=0}^{\infty} h^j M_j f.$$

Star products are the object of study of *deformation quantization*, and were first introduced in the seminal paper by Bayen, Flato, Fronsdal, Lichnerowicz and Sternheimer [BF]. Some more information about them can be found e.g. in the recent surveys by Gutt [Gu] or by S.-T. Ali and the first author [AE].

Toeplitz star product. One can sometimes construct a star product using Toeplitz operators. Namely, under suitable hypothesis on Ω and the Poisson structure (the boundary behaviour of the B_{jk} in (1)), there exists a family of measures μ_h on Ω , $0 < h < 1$, such that the following holds. Let $L_{\text{hol}}^2(\mu_h)$ be the subspace of holomorphic functions in $L^2(\mu_h)$ (weighted Bergman space), $P_h : L^2(\mu_h) \rightarrow L_{\text{hol}}^2(\mu_h)$ the orthogonal projection, and for f a bounded continuous function on Ω define the Toeplitz operator $T_f^{(h)}$ on $L_{\text{hol}}^2(\mu_h)$ by $T_f^{(h)}\phi = P_h(f\phi)$. Then for any $f, g \in \mathcal{D}(\Omega)$ (:=the functions in $C^\infty(\Omega)$ with compact support), there is an asymptotic expansion

$$(5) \quad T_f^{(h)} T_g^{(h)} \simeq \sum_{j=0}^{\infty} h^j T_{C_j(f,g)}^{(h)} \quad \text{as } h \rightarrow 0,$$

with some bidifferential operators C_j (independent of f, g). Further, these operators C_j define — via the formula (2) — a star product on Ω .

Here the expansion (5) is understood in the sense of operator norms, i.e. for each $N = 0, 1, 2, \dots$,

$$(6) \quad \left\| T_f^{(h)} T_g^{(h)} - \sum_{j=0}^N h^j T_{C_j(f,g)}^{(h)} \right\| \leq C_{N,f,g} h^{N+1} \quad \forall h \in (0, 1).$$

Remark. Formally, we can write (5) as

$$T_f^{(h)} T_g^{(h)} = T_{f *' g}^{(h)}. \quad \square$$

Example. [Cob] If $\Omega = \mathbf{C}$ and $d\mu_h(z) = (\pi h)^{-1} e^{-|z|^2/h} dz$ (where dz stands for the two-dimensional Lebesgue measure), then (5) holds with

$$C_j(f, g) = \frac{1}{j!} \frac{\partial^j f}{\partial z^j} \frac{\partial^j g}{\partial \bar{z}^j}.$$

Similarly, (5) holds for $\Omega = \mathbf{D}$, the unit disc, with the standard weighted Bergman spaces $d\mu_h(z) = \frac{h+1}{\pi h}(1 - |z|^2)^{1/h} dz$ [KL].

Other situations when the Berezin-Toeplitz quantization can be carried out include bounded symmetric domains [BLU] (see also [E3] for an extension from functions in \mathcal{D} to functions not necessarily having compact support), strictly pseudoconvex domains with Poisson brackets having a reasonable boundary behaviour [E1], or, provided one considers the manifold case and allows also spaces L_{hol}^2 of sections of lines bundles (instead of just functions), all compact Kähler manifolds [BMS], [Sch].

The Berezin-Toeplitz quantization prompts the following definition.

Definition. We say that a star product (2) is *induced by operators* if there exists a family of Hilbert spaces \mathcal{H}_h , $0 < h < 1$, a “large” subspace $\mathcal{Z} \subset C^\infty(\Omega)$, and linear maps $f \mapsto Q_f^{(h)}$ from \mathcal{Z} into bounded linear operators on \mathcal{H}_h such that

$$(7) \quad Q_f^{(h)} Q_g^{(h)} \simeq \sum_{j=0}^{\infty} h^j Q_{C_j(f,g)}^{(h)} \quad \text{as } h \rightarrow 0, \quad \forall f, g \in \mathcal{Z},$$

in the sense of operator norms.

Here being “large” can be interpreted, for instance, as follows:

for each finite set of multiindices $\alpha_1, \dots, \alpha_k$, complex numbers w_1, \dots, w_k , and point $z \in \Omega$, there exists $f \in \mathcal{Z}$ such that

$$(8) \quad D^{\alpha_j} f(z) = w_j, \quad \forall j = 1, \dots, k,$$

where D^α denotes the operator of differentiation. The merit of (8) is that it ensures that the knowledge of $C_j(f, g)$ for all $f, g \in \mathcal{Z}$ already determines these C_j uniquely.

Conjecture. *Every star product is induced by operators.*

At the moment, we have no ideas how to attack this conjecture. But we are able to prove at least a weaker result:

Main Theorem. *Every star product equivalent to the Toeplitz star product is induced by operators.*

For the case of bounded symmetric domains and star-products which are invariant with respect to holomorphic automorphisms, this theorem was proved by one of the authors in [E2], using heavily the special machinery of Lie groups available in that setup (above all, the Helgason-Fourier transform and the related theory of invariant differential operators on symmetric spaces). No such thing is available in the general case treated here, and thus we use another approach building on an extension of the classical theorem of Borel to Frechet spaces, due to Colombeau [Col].

The paper is organized as follows. In Section 2, we show how the proof of the theorem can be reduced to the problem of existence of a locally convex space of functions on Ω possessing certain properties. This space is then constructed in Section 3 as an inductive limit of a sequence of Banach spaces. The necessary prerequisites on locally convex spaces can be found e.g. in the books of Jarchow [Ja], Koethe [Koe], Meise and Vogt [MV], or Bonnet and Perez Carreras [BnC]. In the final Section 4, we briefly mention also a non-linear variant of the above quantization procedure, for which a much simpler proof can be given.

2. PLAN OF PROOF OF MAIN THEOREM

Let $*$ denote the Toeplitz star product, and let $*'$ be a star product equivalent to $*$. Let $M_0 = I, M_1, M_2, \dots$ be the differential operators furnishing the equivalence, and let M denote the linear operator on $C^\infty(\Omega)[[h]]$ given by (4). Writing (5) and (7) formally as

$$T_f T_g = T_{f*g}, \quad Q_f Q_g = Q_{f*g},$$

and comparing this with (3), we see that if we could take

$$Q_f^{(h)} := T_{Mf}^{(h)}$$

then we would be done. The problem is that Mf is just a formal power series, which may diverge if one assigns to h some value. So we need to approximate, in some sense, the formal power series M by genuine operators.

Denote by C_j the coefficients (2) of the Toeplitz star product, by C'_j the coefficients of $*'$. Expanding (3) and comparing the expressions at like powers of h on both sides, we see that C_j and C'_j are related by

$$(9) \quad \sum_{j+k=N} M_j C'_k(f, g) = \sum_{j+k+l=N} C_j(M_k f, M_l g), \quad \forall f, g \in C^\infty(\Omega),$$

for each $N = 0, 1, 2, \dots$.

Assume that we can construct a vector space \mathcal{Z} contained in $C^\infty(\Omega)$ such that

- (a) $\mathcal{D} \cap \mathcal{Z}$ is “large”, in the sense of (8);
- (b) $M_j \mathcal{Z} \subset L^\infty \forall j$;
- (c) $C_l(M_j \mathcal{Z}, M_k \mathcal{Z}) \subset L^\infty \forall j, k, l$;
- (d) $C'_k(\mathcal{Z}, \mathcal{Z}) \subset \mathcal{Z} \forall k$; and, finally,
- (e) there exists a family of linear operators $M^{(h)}, 0 < h < 1$, from \mathcal{Z} into L^∞ , such that for each $N = 0, 1, 2, \dots$ and $f \in \mathcal{Z}$,

$$\|(M^{(h)} - \sum_{j=0}^N h^j M_j) f\|_\infty \leq C_{f,N} h^{N+1}$$

with some finite constant $C_{f,N}$, for all $0 < h < 1$.

(Note that, as $M_0 = I$, (b) implies in particular that $\mathcal{Z} \subset L^\infty$.)

Granted this, let us set, for $f \in \mathcal{Z}$,

$$Q_f^{(h)} := T^{(h)}[M^{(h)} f],$$

where, for typographical reasons, we started writing $T^{(h)}[f]$ instead of $T_f^{(h)}$. Since the norm of a Toeplitz operator always satisfies

$$\|T_f^{(h)}\| \leq \|f\|_\infty,$$

we see from (e) that, for each $N = 0, 1, 2, \dots$,

$$(10) \quad \|T^{(h)}[M^{(h)} f] - \sum_{j=0}^N h^j T^{(h)}[M_j f]\| = O(h^{N+1}).$$

(Note that $M^{(h)}f, M_j f \in L^\infty$ in view of (e) and (b).)

If g is another function from \mathcal{Z} , it follows that

$$(11) \quad \|Q_f^{(h)}Q_g^{(h)} - \left(\sum_{j=0}^N h^j T^{(h)}[M_j f]\right)\left(\sum_{j=0}^N h^j T^{(h)}[M_j g]\right)\| = O(h^{N+1}).$$

Finally, if f and g (and, hence, also $M_j f, M_k g$) belong in addition to \mathcal{D} , then (6) applies to $T^{(h)}[M_j f]T^{(h)}[M_k g]$, for each j and k ; thus for any $f, g \in \mathcal{D} \cap \mathcal{Z}$

$$(12) \quad \begin{aligned} Q_f^{(h)}Q_g^{(h)} &= \sum_{j,k=0}^N h^{j+k} T^{(h)}[M_j f]T^{(h)}[M_k g] + O(h^{N+1}) \\ &= \sum_{j,k,l=0}^N h^{j+k+l} T^{(h)}[C_l(M_j f, M_k g)] + O(h^{N+1}) \quad \text{by (6)} \\ &= \sum_{j+k+l \leq N} h^{j+k+l} T^{(h)}[C_l(M_j f, M_k g)] + O(h^{N+1}) \quad \text{by (c)} \\ &= \sum_{j+k \leq N} h^{j+k} T^{(h)}[M_j C'_k(f, g)] + O(h^{N+1}) \quad \text{by (9)} \\ &= \sum_{j,k=0}^N h^{j+k} T^{(h)}[M_j C'_k(f, g)] + O(h^{N+1}) \quad \text{by (d) and (b)} \\ &= \sum_{j=0}^N h^j T^{(h)} \left[M_j \sum_{k=0}^N h^k C'_k(f, g) \right] + O(h^{N+1}) \\ &= T^{(h)} \left[M^{(h)} \sum_{k=0}^N h^k C'_k(f, g) \right] + O(h^{N+1}) \quad \text{by (d) and (10) again} \\ &= Q^{(h)} \left[\sum_{k=0}^N h^k C'_k(f, g) \right] + O(h^{N+1}). \end{aligned}$$

(All the O -terms relate to errors in operator norm.) Thus (7) holds (with C'_j in the place of C_j), and the proof is complete.

It thus only remains to construct the space \mathcal{Z} with the above properties.

3. CONSTRUCTION OF THE SPACE \mathcal{Z}

Recall that the classical theorem of Borel asserts that for any sequence f_n of complex numbers, there exists a function $f \in \mathcal{D}(\mathbf{R})$ such that

$$f^{(j)}(0) = j! f_j, \quad \forall j = 0, 1, 2, \dots$$

In particular, by Taylor's formula, it follows that

$$|f(h) - \sum_{j=0}^N h^j f_j| \leq C_{f,N} h^{N+1}, \quad \forall h \in \mathbf{R}.$$

Comparing this with the condition (e), we see that (e) is tantamount to having a Borel theorem for functions on \mathbf{R} with values in the space of operators from \mathcal{Z} into L^∞ equipped with the strong operator topology (the topology of uniform convergence in norm on finite subsets of \mathcal{Z}). Unfortunately, it turns out that in this generality, i.e. for functions with values in a locally convex space, Borel's theorem may fail in general (see [Col]).

However, it is a notable result of Colombeau [Col] that Borel's theorem is valid for functions on \mathbf{R} with values in a Frechet space. Furthermore, it is known that if \mathcal{Z} an (LB) -space, that is, a countable inductive limit of Banach spaces, then the space $L_b(\mathcal{Z}, L^\infty(\Omega))$ of continuous linear operators, endowed with its natural locally convex topology of uniform convergence on bounded sets, automatically becomes a Frechet space. Consequently, if our \mathcal{Z} will be an (LB) -space, then Colombeau's result applies, and we get our requirement (e) granted.

Our purpose will therefore be to construct an (LB) -space $\mathcal{Z} \subset C^\infty(\Omega)$ which satisfies the conditions (a)–(d).

Let $c_{j\alpha} \in C^\infty(\Omega)$ be the coefficients of the differential operators M_j , i.e.

$$M_j f(x) := \sum_{\alpha \text{ multiindex}} c_{j\alpha}(x) D^\alpha f(x);$$

and similarly denote

$$\begin{aligned} C_j(M_k f, M_l g) &= \sum_{\alpha, \beta} c_{jkl\alpha\beta} D^\alpha f \cdot D^\beta g, \\ C'_k(f, g) &= \sum_{\alpha, \beta} c'_{k\alpha\beta} D^\alpha f \cdot D^\beta g. \end{aligned}$$

Let us enumerate the countable set $\{c_{j\alpha}\}_{j,\alpha} \cup \{c_{jkl\alpha\beta}\}_{j,k,l,\alpha,\beta} \cup \{c'_{k\alpha\beta}\}_{k,\alpha,\beta}$ of all the above coefficients as v_j , $j \in \mathbf{N}$. (Here and in the sequel $\mathbf{N} := \{0, 1, 2, \dots\}$.)

Lemma 1. *There exists a C^∞ weight function $w : \Omega \rightarrow \mathbf{R}^+$ such that $w \geq 1$, $w(x) \rightarrow +\infty$ if x tends to the boundary or to infinity, and*

$$\gamma_{\alpha,j} := \sup_{x \in \Omega} \frac{|D^\alpha v_j(x)|}{w(x)} < \infty$$

for all $j \in \mathbf{N}$ and all multiindices α .

Proof. Enumerate the (countable) set of all the functions $D^\alpha v_j$, $j \in \mathbf{N}$, $\alpha \in \mathbf{N}^n$, as f_k , $k = 1, 2, \dots$. Further, pick a sequence ϕ_j of functions in $\mathcal{D}(\Omega)$ such that $0 \leq \phi_j \leq 1$, the union of the supports of ϕ_j is all of Ω , and $\phi_{j+1} = 1$ on the support of ϕ_j . Denote $K_j := \text{supp } \phi_j$, and set $f_0(x) := \sum_j (1 - \phi_j)$. Thus K_j is an increasing sequence of compact subsets such that the union of their interiors is Ω , and f_0 is in $C^\infty(\Omega)$, $f_0 \geq 0$ and $f_0(x) \rightarrow +\infty$ as x tends to the boundary of Ω or to infinity. Now choose constants C_m such that

$$\sup_{x \in K_m, j \leq m, |\alpha| \leq m} |D^\alpha f_j(x)| \leq C_m.$$

Set

$$w(x) := 1 + \sum_{j=0}^{\infty} \frac{f_j(x)}{2^j C_j}.$$

The sum converges uniformly on compact subsets, together with all its partial derivatives, thus $w \in C^\infty(\Omega)$. Clearly $w \geq 1$, and $w(x) \rightarrow +\infty$ as x tends to the boundary or to infinity since $w \geq C_0^{-1}f_0$. Finally, $f_k/w \leq 2^k C_k$. Thus w does the job we need. \square

We fix a function w as in the last lemma from now on, and also set

$$W(x) := e^{w(x)}.$$

Denote, for $j \in \mathbf{N}$,

$$\kappa_j := \sup_{x \in \Omega} \frac{w(x)^j}{W(x)}.$$

From the fact that $w \rightarrow +\infty$ as x tends to the boundary or to infinity it follows that each κ_j is finite.

Finally, we fix from now on a sequence ϕ_k of functions in \mathcal{D} such that the union of their supports is Ω , and $\phi_{k+1} = 1$ on the support of ϕ_k .

Lemma 2. *There exists a sequence $\{\epsilon_j\}_{j \in \mathbf{N}}$ such that $0 < \epsilon_j \leq 1$ for all j and*

$$\sup_{x \in \Omega} W(x) \sum_{\alpha} \epsilon_{|\alpha|} |D^\alpha(x^\gamma \phi_k(x))| < \infty$$

for any $k \in \mathbf{N}$ and any multiindex γ . Here the summation extends over all multiindices $\alpha \in \mathbf{N}^n$, and $|\alpha| := \alpha_1 + \dots + \alpha_n$.

Proof. Choose again constants $C'_m < \infty$ such that

$$\sup_{x \in \Omega, |\gamma| \leq m, k \leq m, |\alpha| \leq m} W(x) |D^\alpha(x^\gamma \phi_k(x))| \leq C'_m.$$

Set $\epsilon_m := 2^{-m}/C'_m$. Then for any k, γ and x ,

$$\sum_{|\alpha| \geq \max(k, |\gamma|)} W(x) |D^\alpha(x^\gamma \phi_k(x))| \epsilon_{|\alpha|} \leq \sum_{\alpha} 2^{-|\alpha|} = 2^n < \infty,$$

and the claim follows. \square

Lemma 3. *There exist positive constants A_k , $k \in \mathbf{N}$, such that $A_k \geq \gamma_{0,k}$ and*

$$(13) \quad A_k A_{|\iota|} \geq 2^{|\iota|} \binom{\iota}{\mu} \gamma_{\iota-\mu, k} \binom{\mu}{\pi}$$

for any multiindices ι, μ, π such that $\pi \subset \mu \subset \iota$.

Here we are using the usual multiindex notation

$$\binom{\iota}{\mu} := \binom{\iota_1}{\mu_1} \dots \binom{\iota_n}{\mu_n},$$

and $\pi \subset \mu$ means that $\pi_j \leq \mu_j$ for all j .

Proof. The right-hand side of (13) can be bounded by a constant $A(k, \iota)$ depending on k and ι only. Take

$$A_m := \max_{k, |\iota| \leq m} A(k, \iota) + 1.$$

It is easy to see that (13) follows. \square

We now define, inductively, a sequence of small positive numbers $a_{m,i}$, $m, i \in \mathbf{N}$. Assume that m and i are given and that $a_{k,j}$ has already been defined for all (k, j) with $k < m$, or $k = m$ and $j < i$. We choose $a_{m,i} > 0$ so small that the following requirements are satisfied:

$$(14) \quad a_{m,i} \leq \epsilon_i \leq 1,$$

$$(15) \quad a_{m,i} \leq a_{m-1,i} \quad \text{if } m \geq 1,$$

$$(16) \quad a_{m,i} \leq a_{m-1,i+1} \quad \text{if } m \geq 1,$$

and

$$(17) \quad a_{m,i} \leq \frac{a_{m,p} a_{m,q}}{A_i} \quad \text{for all } p, q < i.$$

Lemma 4. For all $k, m \in \mathbf{N}$ and $\mu, \pi \in \mathbf{N}^n$ with $\pi \subset \mu$,

$$\sum_{\iota \supset \mu} a_{m,|\iota|} \binom{\iota}{\mu} \gamma_{\iota-\mu,k} \binom{\mu}{\pi} \leq (2^n + 1) A_k \frac{a_{m,|\mu-\pi|} a_{m,|\pi|}}{a_{m,0}},$$

where A_k is as in (13).

Proof. As a combination of (13) and (17) we obtain

$$\begin{aligned} \sum_{\iota \supset \mu} a_{m,|\iota|} \binom{\iota}{\mu} \gamma_{\iota-\mu,k} \binom{\mu}{\pi} &\leq A_k \sum_{\iota \supset \mu} a_{m,|\iota|} A_{|\iota|} 2^{-|\iota|} \\ &\leq A_k a_{m,|\mu-\pi|} a_{m,|\pi|} \sum_{\iota \supset \mu} 2^{-|\iota|} \\ &\leq 2^n A_k a_{m,|\mu-\pi|} a_{m,|\pi|} \leq 2^n A_k \frac{a_{m,|\mu-\pi|} a_{m,|\pi|}}{a_{m,0}}, \end{aligned}$$

since $a_{m,0} \leq 1$ by construction. The remaining term $\iota = \mu$ reduces to $a_{m,|\mu|} \gamma_{0,k} \binom{\mu}{\pi}$. For $0 < |\pi| < |\mu|$ we can again use (13) and (17) to bound this by $A_k a_{m,|\mu-\pi|} a_{m,|\pi|} 2^{-|\mu|} \leq A_k a_{m,|\mu-\pi|} a_{m,|\pi|}$. For $\pi = 0$ or $\pi = \mu$, the term becomes just $a_{m,|\mu|} \gamma_{0,k}$, which is again bounded by $\frac{a_{m,|\mu|} a_{m,0}}{a_{m,0}} A_k$ since $A_k \geq \gamma_{0,k}$ by Lemma 3. \square

We proceed to define the space $\mathcal{Z} := \text{ind}_{k \rightarrow \infty} \mathcal{Z}_k$. The step spaces \mathcal{Z}_k are defined to consist of C^∞ functions f on Ω such that

$$\|f\|_k := \sup_{x \in \Omega} W(x) w(x)^{-k} \sum_{\alpha} a_{k,|\alpha|} |D^\alpha f(x)| < \infty.$$

We have $\mathcal{Z}_k \subset \mathcal{Z}_{k+1}$ continuously, for all k . In fact, $\|f\|_{k+1} \leq \|f\|_k$ by (15) and the fact that $w \geq 1$.

To recall the basic properties of the inductive limit, we have $\mathcal{Z} := \cup_k \mathcal{Z}_k$, and denoting $U_k := \{f \in C^\infty(\Omega) : \|f\|_k \leq 1\}$, a basis of neighbourhoods of zero is given by the sets

$$(18) \quad U := \Gamma \bigcup_{k=1}^{\infty} b_k U_k := \left\{ \sum_k \lambda_k g_k : \sum_k |\lambda_k| \leq 1, g_k \in b_k U_k \right\}$$

for $b_j > 0$. Here the sum in the definition of U is allowed to have only a finite number of terms.

We are now going to prove that the space \mathcal{Z} satisfies the conditions (a)–(e) from the preceding section. This will finish the proof of our Main Theorem.

Proof of (a). By Lemma 2 and (14), all functions of the form $p\phi_k$, where $k \in \mathbf{N}$ and p is an arbitrary polynomial, belong to \mathcal{Z} . Since these functions obviously also belong to \mathcal{D} and the germ of $p\phi_k$ at any point of $\text{supp } \phi_{k-1}$ coincides with the germ of p at that point, it is obvious that $\mathcal{D} \cap \mathcal{Z}$ is “large” in the sense of (8). \square

Proof of (b). It is enough to show that for each l and α , the operator $v_l D^\alpha$ maps \mathcal{Z} into L^∞ . However, if $f \in \mathcal{Z}_k$, then

$$\begin{aligned} |v_l(x)D^\alpha f(x)| &\leq |v_l(x)| \frac{\|f\|_k w(x)^k}{a_{k,|\alpha|} W(x)} \\ &\leq \gamma_{0,l} \frac{\|f\|_k w(x)^{k+1}}{a_{k,|\alpha|} W(x)} \\ &\leq \gamma_{0,l} \frac{\|f\|_k}{a_{k,|\alpha|}} \kappa_{k+1} < \infty \end{aligned}$$

for any $x \in \Omega$. \square

Proof of (c). Again, it is enough to prove that $v_l(D^\alpha f)(D^\beta g) \in L^\infty$ for any $l \in \mathbf{N}$, $\alpha, \beta \in \mathbf{N}^n$ and $f, g \in \mathcal{Z}$. However, for $f \in \mathcal{Z}_k$ and $g \in \mathcal{Z}_m$, by a similar argument as above,

$$|v_l(D^\alpha f)(D^\beta g)| \leq \gamma_{0,l} w \frac{\|f\|_k w^k}{a_{k,|\alpha|} W} \frac{\|g\|_m w^m}{a_{m,|\beta|} W} \leq \gamma_{0,l} \frac{\|f\|_k \|g\|_m}{a_{k,|\alpha|} a_{m,|\beta|}} \kappa_{k+1} \kappa_m < \infty. \quad \square$$

Proof of (d). Once more, we need only to prove that $v_k(D^\alpha f)(D^\beta g)$ belongs to \mathcal{Z} whenever $k \in \mathbf{N}$, α, β are multiindices and $f, g \in \mathcal{Z}$.

Assume that a neighbourhood of 0 as in (18) is given. So we are given a positive sequence $\{b_j\}_{j \in \mathbf{N}}$.

We pick numbers $\tilde{b}_j > 0$, $j \in \mathbf{N}$, such that for all j we have

$$(19) \quad \tilde{b}_j \leq \min_{m \leq 2j+|\alpha|+|\beta|} \left(\frac{b_m a_{m,0}}{\kappa_{m+1} (2^n + 1) A_k}, 1 \right).$$

We claim that $f \in \Gamma \cup_{j=1}^\infty \tilde{b}_j U_j$ and $g \in \Gamma \cup_{j=1}^\infty \tilde{b}_j U_j$ imply

$$v_k(D^\alpha f)(D^\beta g) \in \Gamma \bigcup_{j=1}^\infty b_j U_j.$$

By the assumptions on f and g , we can find finite sets of complex numbers λ_j and μ_j and functions $f_j \in \tilde{b}_j U_j$ and $g_j \in \tilde{b}_j U_j$ such that

$$\sum_j |\lambda_j| \leq 1, \quad \sum_j |\mu_j| \leq 1, \quad f = \sum_j \lambda_j f_j \quad \text{and} \quad g = \sum_j \mu_j g_j.$$

We have

$$v_k(D^\alpha f)(D^\beta g) = \sum_{j,l} \lambda_j \mu_l v_k(D^\alpha f_j)(D^\beta g_l),$$

and here $\sum_{j,l} |\lambda_j \mu_l| \leq 1$, so it is enough to prove that for all j and l there exists m such that $v_k(D^\alpha f_j)(D^\beta g_l) \in b_m U_m$.

We claim that this happens for $m := j + l + |\alpha| + |\beta|$. Recall that $|D^\ell v_k| \leq \gamma_{\ell,k} w$ by Lemma 1. We have

$$(20) \quad \|v_k(D^\alpha f_j)(D^\beta g_l)\|_m = \sup_{x \in \Omega} W(x) w(x)^{-m} \sum_{\ell} a_{m,|\ell|} |D^\ell(v_k(D^\alpha f_j)(D^\beta g_l))|.$$

By the Leibniz rule,

$$\begin{aligned} \sum_{\ell} a_{m,|\ell|} |D^\ell(v_k(D^\alpha f_j)(D^\beta g_l))| &\leq \sum_{\ell} a_{m,|\ell|} \sum_{\mu \subset \ell} \binom{\ell}{\mu} |(D^{\ell-\mu} v_k) D^\mu((D^\alpha f_j)(D^\beta g_l))| \\ &\leq \sum_{\ell} a_{m,|\ell|} \sum_{\mu \subset \ell} \binom{\ell}{\mu} \gamma_{\ell-\mu,k} w \sum_{\pi \subset \mu} \binom{\mu}{\pi} |(D^{\alpha+\mu-\pi} f_j)(D^{\beta+\pi} g_l)| \\ &= w \sum_{\pi} \sum_{\mu \supset \pi} \sum_{\ell \supset \mu} a_{m,|\ell|} \binom{\ell}{\mu} \gamma_{\ell-\mu,k} \binom{\mu}{\pi} |(D^{\alpha+\mu-\pi} f_j)(D^{\beta+\pi} g_l)|. \end{aligned}$$

By Lemma 4 this is bounded by

$$\frac{(2^n + 1)A_k}{a_{m,0}} w \sum_{\pi} \sum_{\mu \supset \pi} a_{m,|\mu-\pi|} a_{m,|\pi|} |(D^{\alpha+\mu-\pi} f_j)(D^{\beta+\pi} g_l)|.$$

Hence, (20) can be bounded by

$$\begin{aligned} &\frac{(2^n + 1)A_k}{a_{m,0}} \sup_{x \in \Omega} W(x) w(x)^{-m+1} \sum_{\pi} \sum_{\mu \supset \pi} a_{m,|\mu-\pi|} a_{m,|\pi|} |(D^{\alpha+\mu-\pi} f_j)(D^{\beta+\pi} g_l)| \\ &\leq \frac{(2^n + 1)A_k}{a_{m,0}} \left(\sup_{\Omega} W w^{-m} \sum_{\mu} a_{m,|\mu|} |D^{\alpha+\mu} f_j| \right) \\ &\quad \cdot \left(\sup_{\Omega} W w^{-m} \sum_{\mu} a_{m,|\mu|} |D^{\beta+\mu} g_l| \right) \cdot \left(\sup_{\Omega} \frac{w^{m+1}}{W} \right) \\ &\leq \kappa_{m+1} \frac{(2^n + 1)A_k}{a_{m,0}} \left(\sup_{\Omega} W w^{-m+|\alpha|} \sum_{\mu} a_{m-|\alpha|,|\mu+\alpha|} |D^{\alpha+\mu} f_j| \right) \\ &\quad \cdot \left(\sup_{\Omega} W w^{-m+|\beta|} \sum_{\mu} a_{m-|\beta|,|\mu+\beta|} |D^{\beta+\mu} g_l| \right) \quad \text{by (16) and } w \geq 1 \\ &\leq \kappa_{m+1} \frac{(2^n + 1)A_k}{a_{m,0}} \left(\sup_{\Omega} W w^{-m+|\alpha|} \sum_{\gamma} a_{m-|\alpha|,|\gamma|} |D^{\gamma} f_j| \right) \\ &\quad \cdot \left(\sup_{\Omega} W w^{-m+|\beta|} \sum_{\gamma} a_{m-|\beta|,|\gamma|} |D^{\gamma} g_l| \right) \\ &\leq (2^n + 1)A_k \frac{\kappa_{m+1}}{a_{m,0}} \tilde{b}_j \tilde{b}_l \leq b_m. \end{aligned}$$

Here in the penultimate inequality we have used the fact that

$$\sup_{\Omega} W w^{-m+|\alpha|} \sum_{\gamma} a_{m-|\alpha|,|\gamma|} |D^{\gamma} f_j| = \|f_j\|_{m-|\alpha|} \leq \|f_j\|_j \leq \tilde{b}_j,$$

since $m - |\alpha| = j + l + |\beta| \geq j$ (and similarly for g_l); while the very last inequality follows from (19). \square

Proof of (e). Let us first of all remark that $L_b(\mathcal{Z}, L^\infty(\Omega))$ is indeed a Frechet space. Indeed, since every (LB) -space is a (DF) -space (cf. [Koe], part I, §29.5.(5)), and $L_b(E, F)$ is complete if E is (DF) and F is any complete locally convex space (see [Koe], part II, §39.6(6)), the completeness of $L_b(\mathcal{Z}, L^\infty(\Omega))$ follows. As for metrizability, \mathcal{Z} has a fundamental sequence $(B_n)_{n=1}^\infty$ of bounded sets (by construction, or by the fact that it is (DF) , see [Koe], part I, beginning of section 29.3). Hence every bounded set of \mathcal{Z} is contained in some multiple cB_n , and from the definition of the topology of uniform convergence on bounded sets we see that the countably many sets

$$\mathcal{U}_n := \{T \in L_b(\mathcal{Z}, L^\infty) : \|Tf\|_\infty \leq 1 \text{ for all } f \in B_n\}$$

form a basis of neighbourhoods of 0 in $L_b(\mathcal{Z}, L^\infty)$. Thus $L_b(\mathcal{Z}, L^\infty)$ is Frechet.

Let us now show in detail how the norm estimate (e) follows from Colombeau's result. For the various facts from the differential calculus in Banach spaces, we refer to [Cha].

Recall that a continuous mapping $P : Y \rightarrow X$, where Y and X are Banach spaces, is called a (continuous) j -homogeneous polynomial if there exists a continuous symmetric j -linear mapping $\tilde{P} : Y^j \rightarrow X$ such that $P(x) = \tilde{P}(x, \dots, x)$. The definition for locally convex spaces is the same. Given a P as above, the mapping \tilde{P} is unique and can be constructed from P using the polarization formula ([Cha], Theorems 4.6 and 4.7).

For all $j \in \mathbf{N}$ let us denote by N_j the j -homogeneous polynomial

$$N_j(h) := h^j M_j$$

from \mathbf{R} into $L_b(\mathcal{Z}, L^\infty(\Omega)) =: X$. Let \tilde{N}_j be the corresponding symmetric j -linear mapping. Clearly, both N_j and \tilde{N}_j are continuous.

By the main theorem of [Col] we find a C^∞ -mapping $M : \mathbf{R} \rightarrow X$ such that the n -th derivative of M coincides with \tilde{N}_j . By the explanation after the main theorem in [Col], the j -th derivative of M is even a continuous j -linear mapping from \mathbf{R} to X_B , where $X_B \subset X$ denotes the linear span of some bounded convex balanced subset $B \subset X$. Also, X_B is a Banach space when endowed with the norm $\|x\|_B := 1/\sup\{r > 0 : rx \in B\}$, and the canonical injection $X_B \rightarrow X$ is continuous.

Hence, also $\tilde{N}_j : \mathbf{R}^j \rightarrow X_B$ is continuous. By the Taylor formula for Banach spaces ([Cha], Theorem 8.9), we obtain the error estimate ($M^{(h)} = M(h)$)

$$(21) \quad \left\| M^{(h)} - \sum_{j=0}^N h^j M_j \right\|_B \leq C_N h^{N+1}.$$

Now every bounded set $B \subset X$ is contained in the set

$$(22) \quad U^\circ := \{T \in X : \|Tf\|_\infty \leq 1 \text{ for all } f \in U\}$$

for some neighbourhood of zero U of \mathcal{Z} . Hence,

$$(23) \quad \left\| M^{(h)} - \sum_{j=0}^N h^j M_j \right\|_B \geq \sup_{f \in U} \left\| \left(M^{(h)} - \sum_{j=0}^N h^j M_j \right) f \right\|_\infty.$$

Let now $f \in \mathcal{Z}$ be given. Since U is a neighbourhood of 0, it absorbs f , i.e. there exists a $c > 0$ such that $f \in cU$. The desired norm estimate (e) (for some f -dependent constant) follows from this, (21) and (23). \square

This completes the proof of the Main Theorem.

Remark. Note that in the above proofs we have actually established assertions slightly stronger than the properties (b)–(d), namely that the inclusions given there are in fact continuous: that is, for each j, k and l , M_j maps \mathcal{Z} continuously into L^∞ , $C_l(M_j \cdot, M_k \cdot)$ maps $\mathcal{Z} \times \mathcal{Z}$ continuously into L^∞ , and C'_l even maps $\mathcal{Z} \times \mathcal{Z}$ continuously into \mathcal{Z} . (Well, for (b) and (c), we have only shown that the operators map bounded sets into balls, but this implies continuity by a simple argument: one forms a small enough neighbourhood U of 0 in \mathcal{Z} using just the definition of the inductive limit topology — that is, U is chosen as the balanced convex hull of the union of countably many bounded sets (the unit balls of the step spaces) multiplied by small constants; if the constants are small enough, U is still mapped into the unit ball of L^∞ .) However, this extra piece of information seems to have no implications for the applications to quantization.

4. A NONLINEAR VARIANT

Note that the argument (12) still works even if the operators $M^{(h)}$ and $Q^{(h)}$ are not assumed to be linear, i.e. if we just require that for each fixed $f \in \mathcal{Z}$ there be some functions $M^{(h)}f \in L^\infty$, $0 < h < 1$, such that (e) holds. In that case, the sought operators $M^{(h)}$ can be constructed for $\mathcal{Z} = \mathcal{D}$ directly along the lines of the usual proof of Borel's theorem. Let us include a proof of this assertion for completeness.

Take $\mathcal{Z} = \mathcal{D}(\Omega)$, the subspace of functions in $C^\infty(\Omega)$ with compact support. Then clearly (a)–(d) hold; let us settle (e) (allowing $M^{(h)}$ to be nonlinear). So let $f \in \mathcal{D}$, and denote for brevity $m_j = M_j f \in \mathcal{D}$.

Fix a function $\phi \in C^\infty(\mathbf{R})$ such that $0 \leq \phi \leq 1$ and $\phi(x) = 1$ for $|x| \leq \frac{1}{2}$, $\phi(x) = 0$ for $|x| \geq 1$. Let $\nu_{n,\alpha} := \|D^\alpha m_n\|_\infty$ and set $q_j := \max_{|\alpha|, n \leq j} \nu_{\alpha,n} + 1$, so that $\nu_{n,\alpha} \leq q_n q_{|\alpha|}$. Let now $c_n := \max(2, n!q_n)$ and define

$$u_n(x, h) = \phi(c_n h) m_n(x) h^n.$$

Clearly each u_n is a bounded smooth function on $\Omega \times \mathbf{R}$. Observe that u_n is nonzero only if

$$(24) \quad |h| \leq \frac{1}{c_n} \leq \frac{1}{2},$$

owing to the fact that $c_n \geq 2$. Using the Leibniz rule, we have for any integer $k \geq 0$ and multiindex α ,

$$D_h^k D_x^\alpha u_n(x, h) = D^\alpha m_n(x) \cdot \sum_{j=0}^k \binom{k}{j} D^{k-j} h^n \cdot c_n^j \phi^{(j)}(c_n h),$$

where the subscripts at D indicated the differentiated variable. The last factor on the right-hand side is bounded by $|h|^{-j} s_j$ where $s_j := \sup_{x \in \mathbf{R}} |x^j \phi^{(j)}(x)| < \infty$. The second factor from the right is equal to $\frac{n!}{(n-k+j)!} h^{n-k+j}$, which is bounded by $n! |h|^{n-k+j}$ (and vanishes for $j < k - n$). Thus

$$\begin{aligned} |D_h^k D_x^\alpha u_n(x, h)| &\leq \|D^\alpha m_n\|_\infty \sum_{j=0}^k \binom{k}{j} n! |h|^{n-k} s_j \\ &\leq q_n q_{|\alpha|} n! |h|^{n-k} W_k, \end{aligned}$$

where $W_k := \sum_{j=0}^k \binom{k}{j} s_j < \infty$. Since $c_n \geq n!q_n$, it follows that

$$\begin{aligned} \sum_{n=k+1}^{\infty} |D_h^k D_x^\alpha u_n(x, h)| &\leq q_{|\alpha|} W_k \sum_{n=k+1}^{\infty} q_n n! |h|^{n-k} \\ &\leq q_{|\alpha|} W_k \sum_{n=k+1}^{\infty} \frac{q_n n!}{c_n} |h|^{n-k-1} \quad \text{by (24)} \\ &\leq 2q_{|\alpha|} W_k < \infty. \end{aligned}$$

As α and k can be arbitrary, we see that the series

$$(25) \quad u(x, h) := \sum_{n=0}^{\infty} u_n(x, h)$$

converges in the C^∞ topology to a function $u \in C^\infty(\Omega \times \mathbf{R})$. Further, as each u_n is, in view of (24), supported in $\text{supp } m_n \times [-\frac{1}{2}, \frac{1}{2}] \subset \text{supp } f \times [-\frac{1}{2}, \frac{1}{2}]$, we even have $u \in \mathcal{D}(\Omega \times (-1, 1))$. Since

$$D_h^j u_n(x, 0) = \begin{cases} 0 & \text{if } j \neq n, \\ j! m_j(x) & \text{if } j = n, \end{cases}$$

the C^∞ convergence of (25) implies that $D_h^j u(x, 0) = j! m_j(x)$. By the Taylor remainder formula, we therefore have for any integer $N \geq 0$

$$u(x, h) - \sum_{j=0}^N m_j(x) h^j = \frac{h^{N+1}}{(N+1)!} D_h^{N+1} u(x, \theta(x, h)h)$$

for some $0 \leq \theta(x, h) \leq 1$; consequently,

$$h^{-N-1} \left| u(x, h) - \sum_{j=0}^N m_j(x) h^j \right| \leq \frac{1}{(N+1)!} \|D_h^{N+1} u\|_\infty < \infty, \quad \forall x, h,$$

since u is compactly supported. Thus the choice

$$M^{(h)} f(x) := u(x, h)$$

will do the job we need. \square

Remark. In effect, the above argument proves Borel's theorem for functions from \mathbf{R} into \mathcal{D} , for the case that the prescribed derivatives m_j have supports in a fixed compact set. Perhaps there is some hope that Borel's theorem might hold even for functions from \mathbf{R} into more general locally convex spaces provided some additional hypothesis is placed on the sequence of the prescribed derivatives — for instance, if the target space is a space of operators, when M_j are “tame” in the sense that there exists a shrinking collection U_k of neighbourhoods of zero such that each M_j maps U_{k+1} into U_k , for all k . Note that having larger and larger supports is also the idea behind Colombeau's counterexample [Col] showing that Borel's theorem fails for functions from \mathbf{R} into \mathcal{D} . \square

It should be remarked that from the point of view of physics, the nonlinear quantization treated in this section is probably a rather doubtful business.

REFERENCES

- [AE] S.T. Ali, M. Engliš, *Quantization methods: a guide for physicists and analysts*, Rev. Math. Phys. **17** (2005), 391-490.
- [BF] F. Bayen, M. Flato, C. Fronsdal, A. Lichnerowicz, D. Sternheimer, *Deformation theory and quantization*, Lett. Math. Phys. **1** (1977), 521–530; Ann. Phys. **111** (1978), 61–110 (part I), 111-151 (part II).
- [BnC] J. Bonnet, P. Perez Carreras, *Barreled locally convex spaces*, North-Holland Mathematics Studies 131, Amsterdam, 1987.
- [BMS] M. Bordemann, E. Meinrenken, M. Schlichenmaier, *Toeplitz quantization of Kähler manifolds and $gl(n)$, $n \rightarrow \infty$ limits*, Comm. Math. Phys. **165** (1994), 281–296.
- [BLU] D. Borthwick, A. Lesniewski, H. Upmeyer, *Non-perturbative deformation quantization on Cartan domains*, J. Funct. Anal. **113** (1993), 153–176.
- [Cha] S.B. Chae, *Holomorphy and calculus in normed spaces*, Marcel Dekker, New York-Basel, 1985.
- [Cob] L.A. Coburn, *Berezin-Toeplitz quantization*, Algebraic Methods in Operator Theory, Birkhäuser, 1994, pp. 101–108.
- [Col] J.F. Colombeau, *Infinite dimensional C^∞ mappings with a given sequence of derivatives at a given point*, J. Math. Anal. Appl. **71** (1979), 95–104.
- [E1] M. Engliš, *Weighted Bergman kernels and quantization*, Comm. Math. Phys. **227** (2002), 211-241.
- [E2] M. Engliš, *Berezin-Toeplitz quantization and invariant symbolic calculi*, Lett. Math. Phys. **65** (2003), 59–74.
- [E3] M. Engliš, *Berezin-Toeplitz quantization on the Schwartz space of bounded symmetric domains*, J. Lie Theory **15** (2005), 27–50.
- [Gu] S. Gutt, *Variations on deformation quantization*, preprint math.QA/0003107, Conférence Moshé Flato (Dijon, 1999), vol. I, Math. Phys. Stud. 21, Kluwer, Dordrecht, 2000, pp. 217–254.
- [Ja] H. Jarchow, *Locally convex spaces*, Teubner, Stuttgart, 1981.
- [KL] S. Klimek, A. Lesniewski, *Quantum Riemann surfaces, I: The unit disc*, Comm. Math. Phys. **146** (1992), 103–122.
- [Koe] G. Koethe, *Topological vector spaces I,II*, Springer, Berlin-New York, 1969,1979.
- [MV] R. Meise, D. Vogt, *Introduction to Functional Analysis*, Oxford Univ. Press, New York, 1997.
- [Sch] M. Schlichenmaier, *Deformation quantization of compact Kähler manifolds by Berezin-Toeplitz quantization*, preprint math.QA/9910137, Conference Moshé Flato 1999 (September 1999, Dijon, France), Vol. 2, Kluwer, 2000, pp. 289–306.

MATHEMATICS INSTITUTE, ŽITNÁ 25, 11567 PRAGUE 1, CZECH REPUBLIC and MATHEMATICS INSTITUTE,
 NA RYBNÍČKU 1, 74601 OPAVA, CZECH REPUBLIC
E-mail address: englis@math.cas.cz

UNIVERSITY OF HELSINKI, DEPARTMENT OF MATHEMATICS AND STATISTICS, P.O.Box 68, FIN-00014
 HELSINKI, FINLAND
E-mail address: jari.taskinen@helsinki.fi