

Iterative solution methods for inverse problems: I Regularization methods for linear problems

Barbara Kaltenbacher, University of Graz

22. Juni 2010

Overview

Inverse problems as operator equations

Linear Problems

Compact operators and singular system

Generalized inverse and ill-posedness

A class of regularization methods

- Error representation

- Some examples of methods

- Convergence

- Convergence rates

Inverse problems as operator equations

- ▶ Often, inverse problems can be formulated as operator equations

$$F(x) = y, \quad (1)$$

where $F : \mathcal{D}(F) \rightarrow \mathcal{Y}$ with domain $\mathcal{D}(F) \subset \mathcal{X}$,
 \mathcal{X}, \mathcal{Y} Hilbert spaces.

- ▶ Measurements are usually contaminated with noise, therefore, we assume that noisy data y^δ with

$$\|y^\delta - y\| \leq \delta. \quad (2)$$

are given.

- ▶ Example: EIT: $F : a \mapsto \Lambda_a$, where Λ_a is the Dirichlet-Neumann operator for

$$\nabla(a\nabla u) = 0 \quad \text{in } \Omega$$

Linear Problems

We consider an operator equation

$$Tx = y \quad (3)$$

where $T \in L(X, Y)$ and X and Y are Hilbert spaces.

$\mathcal{R}(T) \subseteq Y$... range of T

$\mathcal{N}(T) \subseteq X$... nullspace of T

$$Q = \text{Proj}_{\overline{\mathcal{R}(T)}}, \quad P = \text{Proj}_{\mathcal{N}(T)},$$

\perp ... orthogonal complement of linear subspace $M \subseteq Z$:

$$M^\perp = \{z \in Z \mid \langle z, m \rangle_Z = 0 \quad \forall m \in M\}$$

T^* ... adjoint operator

$$\langle Tx, y \rangle_Y = \langle x, T^*y \rangle_X \quad \forall x \in X, y \in Y$$

Compact operators and singular system

Theorem

A compact operator $T \in L(X, Y)$ has a singular system $(\sigma_j; u_j, v_j)_{j \in \mathbb{N}}$:
 $(u_j)_{j \in \mathbb{N}} \subseteq X$ and $(v_j)_{j \in \mathbb{N}} \subseteq Y$ orthonormal systems

$$\begin{aligned} Tu_j &= \sigma_j v_j, & \text{span}(u_j)_{j \in \mathbb{N}} &= \mathcal{N}(T)^\perp = \overline{\mathcal{R}(T^*)}, \\ T^* v_j &= \sigma_j u_j, & \text{span}(v_j)_{j \in \mathbb{N}} &= \overline{\mathcal{R}(T)} = \mathcal{N}(T^*)^\perp, \\ \sigma_j &\rightarrow 0 \text{ as } j \rightarrow \infty. \end{aligned} \quad (4)$$

$$Tx = \sum_{j=1}^{\infty} \sigma_j \langle x, u_j \rangle v_j, \quad T^* y = \sum_{j=1}^{\infty} \sigma_j \langle y, v_j \rangle u_j. \quad (5)$$

$$x = \sum_{j=1}^{\infty} \langle x, u_j \rangle u_j + Px, \quad y = \sum_{j=1}^{\infty} \langle y, v_j \rangle v_j + (I - Q)y.$$

Generalized inverse and ill-posedness

T^\dagger ... generalized inverse of T :

$$\forall y \in \mathcal{D}(T^\dagger) = \mathcal{R}(T) + \mathcal{R}(T)^\perp : \quad T^\dagger y = (T|_{\mathcal{N}(T) \rightarrow \mathcal{R}(T)})^{-1} Qy .$$

Compact T with singular system $(\sigma_j; u_j, v_j)_{j \in \mathbb{N}}$:

$$T^\dagger y = \sum_{j=1}^{\infty} \frac{1}{\sigma_j} \langle y, v_j \rangle u_j ,$$

provided this sum converges:

$$y \in \mathcal{D}(T^\dagger) \iff \sum_{j=1}^{\infty} \frac{\langle y, v_j \rangle^2}{\sigma_j^2} < \infty \quad \text{Picard criterion} \quad (6)$$

Note that in general only $\sum_{j=1}^{\infty} \langle y^\delta, v_j \rangle^2 + \|(I - Q)y\|^2 < \infty$ and on the other hand $\sigma_j \rightarrow 0$ as $j \rightarrow \infty$. \rightsquigarrow **ill-posedness**:

Noise in the j th generalized Fourier coefficient $\langle y, v_j \rangle$ is amplified by $\frac{1}{\sigma_j} \rightsquigarrow$ stronger amplification of high frequent noise.

An Example (1-d source identification):

$$\begin{aligned} -\Delta u &= f && \text{in } \Omega \\ u &= 0 && \text{on } \partial\Omega \end{aligned}$$

Identify $f \in L^2(\Omega)$ from given measurements of u

\equiv solve $TX = y$ with $y = u$, $x = f$, $T = \Delta^{-1}$

$X = L^2(\Omega)$, $Y = L^2(\Omega)$ (measure values but not derivatives!)

$\mathcal{R}(T) \subseteq H^2(\Omega) \hookrightarrow L^2(\Omega) \Rightarrow T$ compact.

1-d case $\Omega = (0, 1)$: singular system $((\pi j)^{-2}; \sin(\pi j \cdot), \sin(\pi j \cdot))$,

$$y \in \mathcal{R}(T) + \mathcal{R}(T)^\perp \iff \sum_{j=1}^{\infty} j^4 \left(\int_{\Omega} y(\xi) \sin(\pi j \xi) d\xi \right)^2 < \infty$$

A class of regularization methods: Definition

$$R_\alpha y^\delta := q_\alpha(T^* T) T^* y^\delta \quad (7)$$

$q_\alpha \in C([0, \|T^* T\|])$ depending on some regularization parameter $\alpha > 0$.

Definition of $f(A)$ by spectral theory for

f ... piecewise continuous function

A ... selfadjoint nonnegative definite operator.

Case A compact with eigensystem $(\sigma_j^2; u_j)_{j \in \mathbb{N}}$:

$$f(A)x = \sum_{j=1}^{\infty} f(\sigma_j^2) \langle x, u_j \rangle u_j.$$

Notation: $x_\alpha := R_\alpha y$, $x_\alpha^\delta := R_\alpha y^\delta$, $x^\dagger := T^\dagger y$

A class of regularization methods: Error representation

Reconstruction error for exact data:

$$x^\dagger - x_\alpha = (I - q_\alpha(T^*T)T^*T)x^\dagger = r_\alpha(T^*T)x^\dagger \quad (8)$$

$$r_\alpha(\lambda) := 1 - \lambda q_\alpha(\lambda), \quad \lambda \in [0, \|T^*T\|]. \quad (9)$$

Total error:

$$x^\dagger - x_\alpha^\delta = \underbrace{r_\alpha(T^*T)x^\dagger}_{\text{approximation error}} + \underbrace{R_\alpha(y - y^\delta)}_{\text{propagated noise}}$$

A class of regularization methods: Some examples

- ▶ Tikhonov regularization (Tikh)

$\min \{ \|Tx - y^\delta\|^2 + \alpha \|x - x_0\|^2 \}$, which is equivalent to

$$x_\alpha^\delta = (T^*T + \alpha I)^{-1}(T^*y^\delta + \alpha x_0). \quad (10)$$

- ▶ iterated Tikhonov regularization (iTikh)

$$x_{\alpha,0}^\delta := 0 \quad (11a)$$

$$x_{\alpha,n+1}^\delta := (T^*T + \alpha I)^{-1}(T^*y^\delta + \alpha x_{\alpha,n}^\delta), \quad n \geq 0 \quad (11b)$$

- ▶ Landweber iteration (LW): with $\|T\|^2 \leq 2$ (scaling) $\alpha = \frac{1}{n}$

$$x_0 = 0 \quad (12a)$$

$$x_{n+1} = x_n - T^*(Tx_n - y^\delta), \quad n \geq 0, \quad (12b)$$

Tikh	$q_\alpha(\lambda) = \frac{1}{\lambda + \alpha}$	$r_\alpha(\lambda) = \frac{\alpha}{\lambda + \alpha}$
------	--	---

itTikh	$q_\alpha(\lambda) = \frac{(\lambda + \alpha)^n - \alpha^n}{\lambda(\lambda + \alpha)^n}$	$r_\alpha(\lambda) = \left(\frac{\alpha}{\lambda + \alpha}\right)^n$
--------	---	--

LW	$q_n(\lambda) = \sum_{j=0}^{n-1} (1 - \lambda)^j$	$r_n(\lambda) = (1 - \lambda)^n$
----	---	----------------------------------

TSVD	$q_\alpha(\lambda) = \begin{cases} \lambda^{-1}, & \lambda \geq \alpha \\ 0, & \lambda < \alpha \end{cases}$	$r_\alpha(\lambda) = \begin{cases} 0, & \lambda \geq \alpha \\ 1, & \lambda < \alpha \end{cases}$
------	--	---

A class of regularization methods: Convergence

In all these examples the functions r_α , q_α satisfy

$$\lim_{\alpha \rightarrow 0} r_\alpha(\lambda) = \begin{cases} 0, & \lambda > 0 \\ 1, & \lambda = 0 \end{cases} \quad (13)$$

$$|r_\alpha(\lambda)| \leq C_r \quad \text{for } \lambda \in [0, \|T^* T\|] \quad (14)$$

$$|q_\alpha(\lambda)| \leq \frac{C_q}{\alpha} \quad \text{for } \lambda \in [0, \|T^* T\|] \quad (15)$$

Theorem

If (13) and (14) hold true, then the operators R_α defined by (7) converge pointwise to T^\dagger on $\mathcal{D}(T^\dagger)$ as $\alpha \rightarrow 0$. With the additional assumption (15) the norm of the regularization operators can be estimated by

$$\|R_\alpha\| \leq \sqrt{\frac{(C_r + 1)C_q}{\alpha}}. \quad (16)$$

If $\bar{\alpha}(\delta, y^\delta)$ is a parameter choice rule satisfying

$$\bar{\alpha}(\delta, y^\delta) \rightarrow 0, \quad \text{and} \quad \delta / \sqrt{\bar{\alpha}(\delta, y^\delta)} \rightarrow 0 \quad \text{as } \delta \rightarrow 0, \quad (17)$$

then $(R_\alpha, \bar{\alpha})$ is a regularization method in the sense that

$$\limsup_{\delta \rightarrow 0} \left\{ \|R_{\bar{\alpha}(\delta, y^\delta)} y^\delta - T^\dagger y\| : y^\delta \in Y, \|y^\delta - y\| \leq \delta \right\} = 0 \quad (18)$$

for all $y \in \mathcal{D}(T^\dagger)$.

Convergence rates under source conditions

source wise representation condition

$$x^\dagger = (T^* T)^\mu w, \quad w \in X, \|w\| \leq \rho. \quad (19)$$

T ... smoothing operator \Rightarrow (19) is abstract smoothness condition.

For the above methods (Tikh, itTikh, LW, TSVD), there exist $\mu_0 \in (0, \infty]$ (*qualification*), $C_\mu > 0$ such that

$$\sup_{\lambda \in [0, \|T^* T\|]} |\lambda^\mu r_\alpha(\lambda)| \leq C_\mu \alpha^\mu \quad \text{for } 0 \leq \mu \leq \mu_0. \quad (20)$$

Theorem

Assume that (19) and (20) hold. Then the approximation error and its image under T satisfy

$$\begin{aligned} \|x^\dagger - x_\alpha\| &\leq C_\mu \alpha^\mu \rho, & \text{for } 0 \leq \mu \leq \mu_0, \\ \|Tx^\dagger - Tx_\alpha\| &\leq C_{\mu+1/2} \alpha^{\mu+1/2} \rho, & \text{for } 0 \leq \mu \leq \mu_0 - \frac{1}{2}. \end{aligned}$$

If the regularization parameter α is chosen according to

$$\alpha^{\mu+\frac{1}{2}} \sim \delta \tag{21}$$

then the optimal convergence rate

$$\|x_\alpha^\delta - x^\dagger\| \leq \tilde{C}_\mu \delta^{\frac{2\mu}{2\mu+1}} \text{ for } 0 \leq \mu \leq \mu_0 \tag{22}$$

is obtained.

Remarks

- ▶ a posteriori regularization parameter choice rules (“ μ -free”)
 - ▶ Morozov’s discrepancy principle:
 - $\alpha = \max s.t. \|Tx_\alpha^\delta - y^\delta\| \leq \delta$;
 - optimal rates (24) for $\mu \leq \mu_0 - \frac{1}{2}$ [Morozov 1968];
 - mod.vers.: (24) for $\mu \leq \mu_0$: [Raus 1988, Engl&Gfrerer 1988]
 - ▶ balancing principle (or Lepskii rule) [Goldenshluger&Perverzev 2000, Bauer&Perverzev 2005]:
 - optimal rates (24), also stochastic setting
 - ▶ generalized cross validation [Wahba 1977, Lukas 2006] for stochastic setting
 - ▶ L-curve [Hansen 1992] “ δ -free” (Bakushinski - veto)
- ▶ logarithmic source conditions for severely ill-posed problems [Hohage 1999]
- ▶ alternative choice of regularization term in Tikhonov: TV, L^1 to enhance sparsity \rightsquigarrow analysis in Banach spaces [Burger&Osher 2004, Schöpfer&Louis&Schuster 2006]